

ON THE ISOMORPHISM OF DISCRETE SUBGROUPS OF $SL(2, \mathbb{R})$

Alladi Sitaram

The purpose of this paper is to give a sufficient condition for the isomorphism of two discrete subgroups Γ and Γ' of $SL(2, \mathbb{R})$ in terms of the norms of the primitive hyperbolic elements of Γ and Γ' . The proof exploits some well-known properties of the Selberg zeta function [3].

1. INTRODUCTION

Let G be the group $SL(2, \mathbb{R})$ (that is, the group of two-by-two real matrices of determinant 1). Let Γ be a discrete subgroup of G such that $\Gamma \backslash G$ is compact. We shall also assume that Γ contains no elements of finite order. It is known (see [1, p. 11]) that under these assumptions Γ contains only hyperbolic elements. (An element $\gamma \in \Gamma$ is said to be *hyperbolic* if it has distinct, real eigenvalues.) An element $\gamma \in \Gamma$ is said to be *primitive* if it is not a positive power of any other element of Γ ; clearly, each conjugate of γ will also be primitive. Let $\{P_\alpha\}$ ($\alpha = 1, 2, \dots$) be a complete set of representatives of the primitive hyperbolic conjugacy classes of Γ , and let $N\{P_\alpha\}$ be the norm of P_α (if λ_1 and λ_2 are the eigenvalues of P_α , then $N\{P_\alpha\} = [\max(|\lambda_1|, |\lambda_2|)]^2$). We are now in a position to state our theorem:

THEOREM. *Let Γ and Γ' be discrete subgroups of $G (= SL(2, \mathbb{R}))$, without elements of finite order, and such that $\Gamma \backslash G$ and $\Gamma' \backslash G$ are compact. Let n_1, n_2, \dots be the norms of the primitive hyperbolic classes of Γ , where $n_1 < n_2 < \dots$, and let m_i be equal to the number of primitive classes whose norm is exactly n_i . Let n'_i and m'_i be similarly defined for Γ' . Then, if $n_i = n'_i$ and $m_i = m'_i$ for all i , the subgroups Γ and Γ' are isomorphic (as abstract groups).*

(Note. It is known [2], [4] that $N\{P_\alpha\}_{\alpha=1}^\infty$ has no accumulation point, and therefore we can write the norms of the hyperbolic elements in the order of increasing magnitude.)

Before giving the proof, we require some more facts. Let

$$H = \{Z; Z \in \mathbb{C} \text{ and } \Im Z > 0\}.$$

Then, since Γ is a subgroup of $SL(2, \mathbb{R})$, Γ acts on the upper half-plane H , and under the assumptions on Γ , $\Gamma \backslash H$ is a compact Riemann surface and Γ is its fundamental group. Let p be the genus of $\Gamma \backslash H$. Then, again under the assumptions on Γ , $p > 1$. In [3], A. Selberg introduces the zeta function

$$Z_\Gamma(s) = \prod_{\alpha} \prod_{n=0}^{\infty} [1 - (N\{P_\alpha\})^{-s-n}].$$

Received October 22, 1975.

The author is currently a Visiting Member at the Tata Institute of Fundamental Research, Bombay.

Michigan Math. J. 23 (1976).