

CORRECTION :
ON THE SPECTRUM OF A HYPONORMAL OPERATOR

(This journal, Vol. 17, 305-309)

TAKASHI YOSHINO

(Received November 29, 1966)

In my paper above indicated, Lemma 1 for Theorem 1 is false. Therefore slight corrections are needed in the proof of Theorem 1. Lemma 1 should be omitted and the sentence "hence $\|T^{-1}y\|^2 \leq \|T^{-2}y\|$ and $\|T^{-1}\|^2 \leq \|T^{-2}\|$. Therefore T^{-1} is normaloid by Lemma 1" in the line 2-4 on p. 306 should be replaced by the following: "hence $\|T^{-1}y\|^2 \leq \|T^{-2}y\|$ for all $y \in H$, $\|y\| = 1$. Being replaced y by $T^{-1}x/\|T^{-1}x\|$, $\|x\| = 1$, we have $\|T^{-2}x\|^2 \leq \|T^{-3}x\| \|T^{-1}x\|$ and $\|T^{-1}x\|^4 \leq \|T^{-2}x\|^2 \leq \|T^{-3}x\| \|T^{-1}x\|$. i.e. $\|T^{-1}x\|^3 \leq \|T^{-3}x\|$ for all $x \in H$, $\|x\| = 1$. Repeating this, we have $\|T^{-1}x\|^n \leq \|T^{-n}x\|$ for all $x \in H$, $\|x\| = 1$. This implies $\|T^{-1}\|^n \leq \|T^{-n}\|$ for all n . Therefore T^{-1} is normaloid."

CORRECTION :
SUMMABILITY METHODS OF BOREL TYPE AND
TAUBERIAN SERIES

(This Journal, Vol. 16 (1964), pp. 209-225)

KYUHEI IKENO

(Received January 9, 1967)

Lemma 2.2 which we used in the paper above mentioned is false and this fact is remarked by B. Kuttner and K. Anjaneyulu. Therefore we must give a correct redefinition of a family $B(a, q(p))$ such that $[C_{pk}]$ satisfies (2. 1), (2. 2), (2. 3) and the condition

$$\sum_{k=0}^{\infty} C_{pk} = 1 + o(p^{-1/2}).$$

This family contains Euler, Taylor, S_a , Valiron, Borel method and all the results except Lemma 2.2 are valid.