

## A remark on Hilbert's Nullstellensatz.

By Harley FLANDERS

(Received Nov. 13, 1953)

In [3] Zariski gave a proof of the Nullstellensatz based on the following lemma.

*Let  $k$  and  $K$  be fields such that  $K=k[x_1, \dots, x_n]$ . Then  $K$  is a finite extension of  $k$ .*

Besides the proof of this lemma that Zariski gave, there is another proof in Artin and Tate [1] and a further proof is indicated in the exercises in Bourbaki [2, p. 87, exercise 4 and p. 106, exercise 12]. It is our purpose to give still another proof of this lemma.

Our proof is based on the following well-known results: (1) If  $o$  is an integral domain with quotient field  $k$ , if  $[K:k]=n$ , and if  $O$  is the set of all elements of  $K$  which are integral over  $o$ , then  $O$  is an integral domain and each element of  $K$  can be written in the form  $A/a$  with  $A$  in  $O$  and  $a$  in  $o$ . (2) The field norm  $N_{K/k}$  is multiplicative. (3) If  $A$  is in  $O$  and  $o$  is integrally closed, then  $N_{K/k}A$  is in  $o$ .

Here is the proof. If each  $x_j$  is algebraic over  $k$ , we are finished. The case in which  $x_1, \dots, x_n$  are independent transcendentals is clearly impossible since the polynomial ring  $k[x_1, \dots, x_n]$  is not a field. If neither of these cases prevails, then we may assume that  $x_1, \dots, x_r$  form a transcendence basis of  $K$  over  $k$  for some  $r$  with  $1 \leq r < n$ , set  $F=k(x_1, \dots, x_r)$ , and have  $K$  a finite extension of  $F$ , with say  $[K:F]=m$ . Let  $o=k[x_1, \dots, x_r]$  which is isomorphic to the polynomial ring  $k[X_1, \dots, X_r]$  over  $k$ . By (1) above, there is an element  $f=f(x_1, \dots, x_r)$  in  $o$  such that for each  $j$ ,  $j=r+1, \dots, n$ ,  $z_j=fx_j$  is integral over  $o$ . We select a non-constant polynomial  $g(X_1, \dots, X_r)$  which is relatively prime to  $f(X_1, \dots, X_r)$  (for example  $g(X)=X_1f(X)+1$ ) and consider  $w=1/g(x_1, \dots, x_r)$ . Since  $w$  is in  $K=k[x_1, \dots, x_n]$ , there is a polynomial  $H(X)$  in  $k[X_1, \dots, X_n]$  such that  $w=H(x_1, \dots, x_n)$ . Multiplying this last relation with a sufficiently high power of  $f$  yields a relation of the form  $f^s w=H_1(x_1, \dots, x_r, z_{r+1}, \dots, z_n)$ , where  $H_1(X_1, \dots, X_r, Z_{r+1}, \dots, Z_n)$  is in