

A remark on G. Glauberman's theorem

By Makoto HAYASHI

(Received Jan. 10, 1974)

(Revised May 13, 1974)

In his paper [2], Glauberman proved an interesting theorem for p -stability. The purpose of this note is to prove an immediate corollary of Glauberman's theorem. We introduce notation which will be used in this note. For a finite group H ; H_p denotes a Sylow p -subgroup of H and $\pi(H:p)$ is the set of primes $\bigcup_P \pi(N_H(P)/C_H(P))$, where P ranges over all p -subgroups of H . Other notations are standard (see [3]).

THEOREM*. *Let G be a finite group. Then the following (i), (ii), (iii) are equivalent.*

- (i) $G \triangleright G_2$.
- (ii) $N_G(ZJ(G_p)) \triangleright N_G(ZJ(G_p))_2$, for every odd prime $p \in \pi(G)$.
- (iii) $2 \notin \pi(N_G(ZJ(G_p)):p)$, for every odd prime $p \in \pi(G)$.

COROLLARY. *Let G be a finite group admitting a fixed-point-free automorphism ϕ of a prime power order p^n . Suppose that the order of $C_G(\phi^{p^n-1})$ is odd. Then G is solvable.*

PROOF OF THE THEOREM. (i) \Rightarrow (ii) and (ii) \Rightarrow (iii) follow immediately from definition. To prove (iii) \Rightarrow (i), we assume by way of contradiction that G_2 is not normal in G . Then we have $2 \in \pi(G:p)$ for some odd prime in $\pi(G)$ by Baer's theorem (see [3] page 105). So the order of $N_G(W)/C_G(W)$ is divisible by 2 for some p -subgroup W of G . But the functor ZJ controls strong fusion in G_p with respect to G , because, otherwise, Corollary 2 of Theorem A of [2] would imply that $2 \in \pi(N_G(ZJ(G_p):p))$, contrary to our assumption (iii). Then for $g \in N_G(W)$, we have $g = cn$, $c \in N_G(W)$, $n \in N_G(ZJ(G_p))$. Then we have $n \in N_{N_G(ZJ(G_p))}(W)$, and n has the same action on W as g . So we have $2 \in \pi(N_G(ZJ(G_p)):p)$. This contradicts (iii), q. e. d.

PROOF OF THE COROLLARY. Let G be a minimal counter example group of the corollary. Then all ϕ -invariant proper subgroups of G are solvable.

*) Originally, to prove this theorem, I did not use Glauberman's theorem, but Thompson's N -group classification theorem. I would like to thank the referee for his several suggestions, including improvement in the proof of the theorem.