

A generalized Lüroth Theorem for curves

By T. T. MOH and W. J. HEINZER*

(Received May 25, 1977)

(Revised April 17, 1978)

Let k be a field. The famous "Lüroth Theorem" asserts that if R is a field with $k \subset R \subseteq k(X)$, then $R = k(Y)$, a simple transcendental extension of k . [5, p.198]. As was proved by Igusa [2], [3], Lüroth's Theorem can be generalized to say that if $X_1, \dots, X_n$ are algebraically independent over k and R is a field of transcendence degree one over k such that $k \subset R \subseteq k(X_1, \dots, X_n)$, then $R = k(Y)$, a simple transcendental extension of k . Related results for the case when R has transcendence degree > 1 over k are given by Zariski [6], Swan [4], and Clemens-Griffiths [1].

These striking results naturally motivate the search for similar phenomena or generalization. For this purpose we use the following notation. If R is a function field of one variable over k , then the degree of irrationality of R over k , $\text{irr}(R) = \min\{[R : k(x)] : x \in R\}$. The classical Lüroth Theorem can then be stated: if $R \subseteq S$ are function fields of one variable over k and $\text{irr}(S) = 1$, then $\text{irr}(R) = 1$. In this form, Lüroth's Theorem naturally calls for the study of the pair of numbers $(\text{irr}(S), \text{irr}(R))$ for the case $\text{irr}(S) > 1$. Our result is the following.

THEOREM. *Let $R \subseteq S$ be function fields of one variable over a field k . For any $x \in S$, let y denote the norm of x with respect to R . If y is not algebraic over k , then $[S : k(x)] \geq [R : k(y)]$. In particular, if k is an infinite field, then the degree of irrationality of R , $\text{irr}(R) \leq \text{irr}(S)$, the degree of irrationality of S .*

PROOF. We first consider the case when S is separable over R . Let T be a normal closure of S over R , and let G be the Galois group of T over R . We recall that if H is the subgroup of G fixing S and $G = g_1 H \cup \dots \cup g_m H$ is a coset decomposition of G with respect to H , then $y = \prod_{i=1}^m g_i(x)$ is the norm of x [7, p.91]. Note that $m = [S : R]$. Since $[T : k(x)] = [T : k(g_i(x))]$ is equal to the degree of the polar divisor of x or $g_i(x)$ in T , and since the

* Both authors were supported partially by NSF.

The authors wish to thank the referee for his suggestion that simplified and improved this article.