

A NOTE ON NORMED RING.

By Naoki KIMURA

1. I. Gelfand has shown in his first paper on Normierte Ringe (Recueil Mathématique, T.9 (51), 1941) that if R satisfies four conditions (α) , (β) , (γ) , and (δ) given below, then R is algebraically isomorphic and topologically homeomorphic to R' with the same three conditions (α) , (β) , (γ) , and (δ') which is strictly stronger than (δ) .

According to his proof, he assumed commutativity of R or, at least, the existence of right unit element of R . In this note, we shall show that his assertion is still valid in the case without assumption such as commutativity of R .

It is to be mentioned, however, that our condition (γ) has a right unit element, while Gelfand's (γ) has a left unit element.

2. Let R be a set of elements x, y, z, \dots which satisfies the following four conditions (α) , (β) , (γ) and (δ) .

(α) R is a Banach space with complex numbers as its coefficient field.

(β) R is a ring:

$$x(\lambda y + \mu z) = \lambda xy + \mu xz$$

(λ, μ are complex numbers),

$$x(yz) = (xy)z$$

(γ) R has a right unit element e :

$$xe = x$$

moreover $ne \neq 0$

(δ) Operation of Multiplication is continuous, i.e.,

$$x_n \rightarrow x \text{ implies } yx_n \rightarrow yx,$$

$$\text{and } x_n y \rightarrow xy.$$

Let Q be a Banach space of all linear operators on R into R itself. And let R' be the totality of A_x in Q such that

$$A_x y = xy,$$

$$\text{i.e., } R' = (A_x; x \in R)$$

Then, for the mapping $\varphi: x \leftrightarrow A_x$ between R and R' , we can easily show that

$$(1) \quad x \neq x' \text{ implies } A_x \neq A_{x'},$$

which evidently asserts a one-to-one mapping of φ between R and R' .

$$(2) \quad \varphi \text{ is algebraic isomorphism.}$$

$$(3) \quad \varphi \text{ is continuous from } R' \text{ onto } R.$$

$$(4) \quad R' \text{ is closed in } Q; \text{ thus } R' \text{ is complete.}$$

Therefore by the known theorem of Banach,

$$(5) \quad \varphi \text{ is continuous from } R \text{ onto } R'.$$

We can then conclude that

R and R' are isomorphic and homeomorphic, and moreover R' satisfies the stronger conditions

$$(\gamma') \quad \|e\| = 1.$$

$$(\delta') \quad \|xy\| \leq \|x\| \cdot \|y\|$$

Proof (1)

If $x \neq x'$, then

$$A_x e = xe = x \neq x' = x'e = A_{x'} e$$

Hence $A_x \neq A_{x'}$

In the case of Gelfand, (1) is not satisfied, and we shall give its counter example at the end of this note (4. (b)).

(2) Obvious.

(3) By the inequality $\|A_x\| \geq \frac{1}{\|x\|} \|x\|$.

(4) If $A_{x_n} \rightarrow A \in Q$, then $\{x_n\}$ is a Cauchy sequence, for

$$\|x_n - x_m\| \leq \|A_n - A_m\| \rightarrow 0$$

($n, m \rightarrow \infty$)

R being complete, there exists an element $x \in R$, such that

$$x_n \rightarrow x \quad (n \rightarrow \infty)$$