

ALEXANDROFF'S MAPPING THEOREM FOR PARACOMPACT SPACES

By Keio NAGAMI

(Comm. by H. Tôyama)

This paper states Alexandroff's mapping theorem for paracompact spaces and gives a new characterization of paracompact spaces. A topological space R is called to be approximated by complexes with geometric, natural or weak topology if, for every open covering \mathcal{U} of R , there exist respectively a simplicial complex K with geometric¹⁾, natural²⁾ or weak topology³⁾ and a continuous mapping $f|R \rightarrow K$ such that $\{f^{-1}(S(p))\}$ refines \mathcal{U} , where $S(p)$ denotes an open star with a centre p and p runs through all vertices of K . C. H. Dowker [1] has proved that every paracompact Hausdorff space is approximated by geometric complexes or by natural ones. Our result (Theorem 1) asserts that every paracompact Hausdorff space is approximated by complexes with weak topology. Since weak topology is weaker than geometric and natural topology, ours includes Dowker's results.

Theorem 1. A paracompact Hausdorff space R is approximated by complexes with weak topology.

Proof. Let \mathcal{U} be an arbitrary open covering of R , and then there exists a locally finite open covering $\mathcal{V} = \{V_\alpha : \alpha \in A\}$ of R which refines \mathcal{U} . Since R is normal and then \mathcal{V} is shrinkable, we can assume with no loss of generality that each V_α is an F_σ -set. Therefore, there is a non-negative real-valued continuous function f_α defined on R such that $f_\alpha(x) > 0$ if and only if $x \in V_\alpha$. We associate with each V_α a mark p_α , and with p_α 's as vertices we construct the nerve K of \mathcal{V} such that p_α, \dots, p_β are vertices of a simplex of K if and only if the corresponding sets V_α, \dots, V_β have a common point. We introduce into K the weak topology. Let x be an arbitrary point of R and $A(x)$ be the set of indices such that $A(x) = \{\alpha : x \in V_\alpha\}$. We construct a

mapping $f|R \rightarrow K$ as follows:

$f(x)$ = the centre of gravity of the vertices of $\{p_\alpha : \alpha \in A(x)\}$ with the weights $f_\alpha(x)$.

Then f is continuous: Let W be an open neighborhood of x such that $B(x) = \{\alpha : W \cap V_\alpha \neq \emptyset\}$ is a finite set of indices. Let K_1 , a subcomplex of K , be the nerve of $\{W \cap V_\alpha : \alpha \in B(x)\}$. Then evidently $f(W) \subset K_1$. Being K_1 a finite complex and f_α continuous, it can easily be seen that $f|W \rightarrow K_1$ is continuous and hence $f|R \rightarrow K$ is continuous. From construction of K , $S(p_\alpha)$ is nothing but the set of all points with a non-zero weight on p_α , and hence f is a baricentric \mathcal{V} -mapping, i. e.

$$f^{-1}(S(p_\alpha)) = V_\alpha \quad \text{for all } \alpha \in A.$$

Thus $\{f^{-1}(S(p_\alpha)) : \alpha \in A\}$ refines \mathcal{U} . Q. E. D.

It is to be noted that a complex K in the above can be reconstructed in more restricted type as follows.

Theorem 2. Each star $S(p)$ of K can be of finite dimension.

Proof. Since a paracompact Hausdorff space is strongly screenable [5], we can assume with no loss of generality that \mathcal{V} stated in the above proof can be decomposed into a sequence \mathcal{V}_i^2 , $i = 1, 2, \dots$, such that:

$$\mathcal{V} = \bigcup_{i=1}^{\infty} \mathcal{V}_i^2$$

$$\mathcal{V}_i^2 = \{V_{\alpha_i} : \alpha_i \in A_i\},$$

V_{α_i} 's are, for fixed i , mutually disjoint.

Setting