

SIXES AND SEVENS

Carl Pomerance

University of Georgia

Cooper and Kennedy [1] have posed the following interesting question. For $n = 0, 1, \dots$, let $a_n \in \{6, 7\}$ be such that the base 10 number $a_{n-1}a_{n-2}\dots a_0$ is divisible by 2^n . It is clear that this condition defines the sequence $\{a_n\}$. The first five terms are 6, 7, 7, 7, 6. Cooper and Kennedy ask if $\{a_n\}$ must contain infinitely many 6's and infinitely many 7's. We show a more general result which immediately answers their question in the affirmative.

Theorem. Let b, c be integers at least 2 and let $\{a_n\}_{n=0}^\infty$ be a sequence with each $a_n \in \{0, 1, \dots, b-1\}$. Suppose that for each positive integer m there is some integer N_m , such that c^m divides $a_0 + a_1b + \dots + a_nb^n$ for each $n \geq N_m$. If $\{a_n\}$ is eventually periodic, then $\{a_n\}$ is identically zero.

Proof. It clearly suffices to prove the result in the case that $c = p$, a prime. First note that the hypothesis that N_m always exists implies that either $\{a_n\}$ is identically zero or $p|b$. Indeed, let

$$A_n := \sum_{i=0}^{n-1} a_i b^i$$

for $n = 1, 2, \dots$. Let m be so large that $p^m \geq b$. If $n > N_m$, then

$$p^m | A_n, \quad p^m | A_{n+1} = A_n + a_n b^n,$$

so that $p^m | a_n b^n$. But $0 \leq a_n \leq b-1$ and $p^m \geq b$. Thus, either $\{a_n\}$ is eventually zero or $p|b$. But if $\{a_n\}$ is eventually zero, then there is some N with $A_n = A_N$ for all $n \geq N$. Thus, $p^m | A_N$ for all m , so that $A_N = 0$ and $\{a_n\}$ is identically zero.