

Yet another application of the Gauss-Bonnet Theorem for the sphere*

J. M. Almira A. Romero

The Gauss-Bonnet Theorem, when applied to any Riemannian metric g on the sphere \mathbb{S}^2 , claims that

$$\int_{\mathbb{S}^2} K dA_g = 4\pi,$$

where K and dA_g denote the Gaussian curvature and the area element of g , respectively. In particular, this implies the existence of at least an elliptic point of (\mathbb{S}^2, g) . In this short note we prove the Fundamental Theorem of Algebra as a consequence of this fact. As far as the authors know, this proof is new. Moreover, it does have the advantage of being directly based on the work of Gauss, who was the first mathematician giving an universally accepted proof of the Fundamental Theorem of Algebra (see [2], [3]). Hence our argument can also have some historical and pedagogical interest.

As it is well known, the stereographic projection allows to identify the sphere \mathbb{S}^2 with the one-point compactification $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ of the complex plane \mathbb{C} . Let us take $p(z) = a_0 + a_1z + \cdots + a_nz^n$ with $a_0 a_n \neq 0$ and let us assume that $p(z) \neq 0$ for all $z \in \mathbb{C}$.

We set $p_*(z) = a_n + a_{n-1}z + \cdots + a_0z^n$, that is, the i -th coefficient of $p_*(z)$ is the $(n - i)$ -th coefficient of $p(z)$. Clearly, $p_*(z) = z^n p(1/z)$ for all $z \in \mathbb{C} \setminus \{0\}$. Put

$$f(z) = p(z) p_*(z),$$

for all $z \in \mathbb{C}$. Then f satisfies the functional equation

$$\left| f\left(\frac{1}{z}\right) \right| = \frac{1}{|z|^{2n}} |f(z)|,$$

*The second author is partially supported by MCYT-FEDER Grant MTM2004-04934-C04-01.

Received by the editors May 2006.

Communicated by L. Vanhecke.