

3. Note on the Fundamental Domain of a General Fuchsian Group.

By Masao SUGAWARA.

Tokyo Bunrika Daigaku, Tokyo.

(Comm. by T. TAKAGI, M.I.A., Jan. 12, 1942.)

Let R be the general Poincaré-space $(Z; N(Z) \leq 1)$ ¹⁾ and $W = (U_1 Z + U_2)(U_3 Z + U_4)^{-1}$ be a displacement of a general fuchsian group \mathfrak{G} ; namely

$$U = \begin{pmatrix} U_1 & U_2 \\ U_3 & U_4 \end{pmatrix}, \quad U' S \bar{U} = S, \quad S = \begin{pmatrix} E_m & 0 \\ 0 & -E_n \end{pmatrix}.$$

Definition. When $U_3 \neq 0$, we denote the domain $(Z; \|U_3 Z + U_4\| \geq 1)$ by the symbol $M(U)$. A point Z is an inner or a boundary or an outer point of $M(U)$ according as $\|U_3 Z + U_4\| > 1$ or $= 1$ or < 1 .

Theorem 1. 1°. If $U_3 \neq 0$, $|E - \bar{W}' W|$ is greater than, equals to, or smaller than $|E - \bar{Z}' Z|$ according as Z is an outer or a boundary or an inner point of $M(U)$; the converse also holds.

2°. If however $U_3 = 0$, $|E - \bar{W}' W| = |E - \bar{Z}' Z|$.

Proof. 1°. It comes from the equality

$$\|U_3 Z + U_4\|^{-2} = |E - \bar{W}' W| |E - \bar{Z}' Z|^{-1},$$

which I deduced in another place²⁾.

2. In this case $W = U_1 Z U_4^{-1}$, where U_1 and U_4 are unitary, so that the result is evident.

We denote the intersection of R and all the domains $M(U)$ by M .

Theorem 2. Let Z_0 be a point which gives the maximum value of $|E - \bar{Z}' Z|$ among the points equivalent to Z_0 and let D be the set of such points as Z_0 , then $D = M$.

Proof. Let Z_1 be an inner point of the space R , the number of such points Z equivalent to Z_1 that give $|E - \bar{Z}' Z| \geq |E - \bar{Z}'_1 Z_1|$ is finite, because the series

$$\sum_{U \in \mathfrak{G}} \|U_3 Z + U_4\|^{-k(n+m)}, \quad k \geq 2^3),$$

is absolutely convergent at an inner point of R .

As a point equivalent to an inner point of R is also an inner point of R , the maximum value of the expression $|E - \bar{Z}' Z|$ is really taken at an inner point of R equivalent to Z_1 .

On the other hand, a point equivalent to a boundary point of R is also a boundary point of R and it is always $|E - \bar{Z}' Z| = 0$, when $N(Z) = 1$. Now let $Z \in M$ and let Y be a point equivalent to Z , namely $Y = (U_1 Z + U_2)(U_3 Z + U_4)$, then $|E - \bar{Z}' Z| \geq |E - \bar{Y}' Y|$, if $U_3 \neq 0$, and

1) $N(Z)$ means the norm of Z .

2) M. Sugawara. On the general Zetafuchsian Functions,

3) In the case of symmetrical matrices the exponent is $-k(n+1)$.