

**100. On the Behaviour of an Inverse Function
of a Meromorphic Function at its Trans-
cendental Singular Point, II.**

By Masatsugu TSUJI.

Tokyo Imperial University.

(Comm. by T. YOSIE, M.I.A., Dec. 12, 1941.)

In my former paper¹⁾, I have proved a theorem concerning the behaviour of an inverse function of a meromorphic function at its transcendental singularity $w=0$.

In the proof of $\lim_{r \rightarrow \infty} A(r) = \infty$, I have tacitly assumed that the boundary of Δ has a branch tending to infinity. I will here give a proof of $\lim_{r \rightarrow \infty} A(r) = \infty$ in the general case, where the boundary of Δ has a branch tending to infinity or it has no such a branch, but it consists of only closed curves. We will follow Mr. K. Kunugui's proof (in the paper cited in my former paper), but simplify it at some points.

Proof of $\lim_{r \rightarrow \infty} A(r) = \infty$.

Case I, where for some w_0 ($0 < |w_0| < \rho$), $w_0 = f(z)$ has only finite number of roots in Δ . Then by Mr. K. Noshiro's extension of Iversen-Valiron's theorem (in the paper cited in my former paper), there exists a curve Γ_1 in Δ tending to infinity, such that $\lim f(z) = w_0$, when z tends to ∞ along Γ_1 . Also, as we have remarked in my former paper, there exists a curve Γ in Δ tending to infinity, such that $\lim f(z) = 0$, when z tends to ∞ along Γ . As before, let $\{a_v^{(r)}\}$ be the part of the boundary of Δ_r which lies on $|z - z_0| = r$ and let $a_v^{(r)}$ intersect Γ . There occurs two cases: (i) two end-points of $a_v^{(r)}$ lie on the boundary of Δ or (ii) the circle $|z - z_0| = r$ does not intersect the boundary of Δ .

In the case (i), we have as before $L(r) \geq \frac{\rho}{2}$ for $r \geq r_1$, and in the case

(ii), since $|z - z_0| = r$ intersects Γ and Γ_1 , $L(r) \geq \frac{|w_0|}{2}$ for $r \geq r_2 > r_1$.

In either case, $L(r) \geq l$ for $r \geq r_2$, where $l = \text{Min.} \left(\frac{\rho}{2}, \frac{|w_0|}{2} \right)$, whence we conclude as before $\lim_{r \rightarrow \infty} A(r) = \infty$.

Case II, where $w = f(z)$ has infinitely many roots in Δ for every w ($0 < |w| < \rho$). Let $n(r, w)$ be the number of roots of $w = f(z)$ in Δ_r , then $\lim_{r \rightarrow \infty} n(r, w) = \infty$.

1) M. Tsuji: On the Behaviour of an Inverse Function of a Meromorphic Function at its Transcendental Singular Point. This Proc. 17 (1941), 414-417. During the correction of my former paper Mr. Y. Tumura's paper: Sur le problème de M. Kunugui appeared in this Proc. 17 (1941), 289-295 and its full detail will appear in the Japanese Journal of Mathematics, 18, where a more general theorem as mine is proved, but in somewhat different definition of $A(r)$ and $S(r)$.