

11. On the Series of Orthogonal Functions.

By Kien-Kwong CHEN.

Mathematical Institute, Tohoku Imp. University, Sendai.

(Rec. Feb. 1, 1928. Comm. by M. FUJIWARA, M.I.A., Feb. 12, 1928.)

Let c_1, c_2, \dots be a sequence of real numbers and $\varphi_1(x), \varphi_2(x), \dots$ a sequence of normalized orthogonal functions in the interval $(0, 1)$; then relating to the series

$$(1) \quad c_1 \varphi_1(x) + c_2 \varphi_2(x) + \dots$$

we have the following theorems :

- (A) If the series $\sum (\log \nu)^2 \cdot c_\nu^2$ is convergent, the series (1) converges almost everywhere in the interval $(0, 1)$ ¹⁾.
 (B) If the series $\sum (\log \log \nu)^2 \cdot c_\nu^2$ is convergent, the series (1) is C_1 -summable almost everywhere in the interval $(0, 1)$ ²⁾.

The author proved that (A) and (B) are deducible from the following theorem, due to Borgen and Kaczmarz²⁾.

- (C) If the series $\sum (\log \log \nu)^2 \cdot c_\nu^2$ is convergent, the sequence of the partial sums

$$(2) \quad s(x, 2), s(x, 2^2), \dots, s(x, 2^n), \dots$$

converges almost everywhere in the interval $(0, 1)$, where

$$s(x, p) = c_1 \varphi_1(x) + c_2 \varphi_2(x) + \dots + c_p \varphi_p(x).$$

Conversely, it can be shown that the theorem (C) is deducible from (A), and also from (B).

Thus, we know that each of the theorems (A) and (B) is deducible from the other ; i. e.

The theorems (A) and (B) are equivalent.

Corresponding to Lebesgue's constants in the theory of trigonometric series, we have for the system of normalized orthogonal functions $\varphi_1(x), \varphi_2(x), \dots$, Lebesgue's functions

1) H. Rademacher, Math. Ann. **87** (1922); D. Menchoff, Fundamenta Mathematicae **4** (1923).

2) D. Menchoff, Comptes Rendus **180** (1925); S. Borgen, Math. Ann. **98** (1927); S. Kaczmarz, Math. Zeits. **26** (1927).