

189. *A Note on a Paper of Farkas*

By Ryutaro HORIUCHI
Kyoto Industry University

(Comm. by Kinjirô KUNUGI, M. J. A., Dec. 12, 1969)

H. E. Rauch [1] and H. M. Farkas [2] discussed analytic submanifolds of Teichmüller space, and in relation to these studies Farkas [3] pointed out the following theorem:

Let S be a compact Riemann surface of genus $g \geq 4$. Let q be a Weierstrass point on S whose Weierstrass sequence begins with 3. Then 4 is a gap at q .

In this paper we shall prove the following more general theorem.

Theorem. *Let S be a compact Riemann surface of genus $g > r(r-1)/2$, where $r(1 < r < g)$ is an integer. Let q be a Weierstrass point on S whose Weierstrass sequence begins with r . Then $r+1$ is a gap at q .*

First we recall some definitions and results from the theory of compact Riemann surfaces.

There are exactly g orders n_i , $0 < n_1 < n_2 < \dots < n_g < 2g$, that can be specified at each point p on S such that no meromorphic function exists having as its only singularity a pole of order n_i at p . The sequence (n_1, n_2, \dots, n_g) is called then a gap sequence at p . Given a point p on S , its gap sequence is $(1, 2, \dots, g)$ in general; however, there do exist points on S whose gap sequences omit some of these numbers. These points are called Weierstrass points. In other words, the gap sequence for a Weierstrass point omits an integer n , $2 \leq n \leq g$. The complement of the gap sequence in the sequence of integers $(1, 2, \dots, 2g)$ is called the Weierstrass sequence.

Lemma. *If there is a Weierstrass point on S whose Weierstrass sequence contains $r, r+1, \dots, r+m$, then*

$$(t+1)[(r-1) - tm/2] \geq g$$

where t is the smallest integer which satisfies $t \geq (r-1)/m$.

Proof. The integers $r, r+1, \dots, r+m$ form the module whose elements are not gaps. Hence the gaps must be contained in the set of remaining natural numbers, which are $1, 2, \dots, r-1; r+m+1, r+m+2, \dots, 2r-1; \dots; \dots, tr-1$; where t is the smallest integer which satisfies $t \geq (r-1)/m$. While as is well known, the number of gaps is exactly g , so we have

$$(r-1) + (r-m-1) + \dots + (r-tm-1) \geq g$$

that is,