

96. Topology of Compact Sasakian Manifolds

By Shûkichi TANNO

Mathematical Institute, Tôhoku University, Sendai

(Comm. by Zyoiti SUETUNA, M.J.A., June 12, 1967)

Introduction. The Betti numbers of compact Kählerian manifolds were studied by many geometers. Especially the second Betti numbers of compact Kählerian manifolds were studied by M. Berger [1], R. L. Bishop and S. I. Goldberg [2] and others. It is natural to research these problems in compact Sasakian manifolds, and in fact, they were studied by S. I. Goldberg [5], S. Tachibana [7], S. Tachibana and Y. Ogawa [8] and S. Tanno [10], etc.

1. In [10] we have used $(m-1)$ -homothetic deformations to get results of the first Betti numbers. We call these deformations D -homothetic deformations, where D denotes the distribution defined by a contact form η . To get results of the second Betti numbers and harmonic forms, we also utilize a D -homothetic deformation:

$$(1.1) \quad g \rightarrow *g = \alpha g + (\alpha^2 - \alpha) \eta \otimes \eta$$

of the associated Riemannian metric g for a positive constant α . Then if (ϕ, ξ, η, g) is a Sasakian structure for a contact form η , then $(*\phi = \phi, *\xi = \alpha^{-1}\xi, *\eta = \alpha\eta, *g)$ is also a Sasakian structure. By studying the relations of harmonic forms with respect to g and $*g$, we get

Theorem 1. *A compact m -dimensional Sasakian manifold M with sectional curvature $> -3/(m-2)$ has the first Betti number $b_1(M) = 0$. If $m = 3$, we have also $b_2(M) = 0$.*

A harmonic 2-form w is called of the hybrid type (pure type, resp.) if it satisfies

$$(1.2) \quad w(\phi X, \phi Y) = w(X, Y) \quad (= -w(X, Y), \text{ resp.})$$

for any vector fields X and Y on M .

Theorem 2. *If $m \geq 5$, a compact Sasakian manifold M with sectional curvature $> -3/(m-2)$ has no harmonic 2-form of the pure type. And if sectional curvature > 0 , there is no harmonic 2-form of the hybrid type. Especially, then, we have $b_2(M) = 0$.*

Remark. Under the additional condition that ξ is regular, S. I. Goldberg [5] obtained the last half of Theorem 2. In Kählerian case the similar fact was obtained by R. L. Bishop and S. I. Goldberg [2].

2. We denote by $K(X, Y)$ the sectional curvature for 2-plane determined by X and Y . As is well known in Kählerian manifolds holomorphic pinchings were studied by several authors. In Sasakian