

## 107. Inverse Images of Closed Mappings. III

By Sitiro HANAI

Osaka University of Liberal Arts and Education

(Comm. by K. KUNUGI, M.J.A., Oct. 12, 1961)

K. Nagami has recently obtained the following theorem:<sup>1)</sup> *a completely regular  $T_1$ -space  $X$  is compact if and only if the projection from the product space  $X \times Y$  onto  $Y$  is closed for any completely regular  $T_1$ -space  $Y$ .*

In this note, with the exception of the complete regularity and the separation axiom ( $T_1$ ) of  $X$ , we shall prove an analogous theorem.

**Theorem.** *Let  $X$  be a topological space and  $m$  an infinite cardinal number. Then  $X$  is  $m$ -compact if and only if the projection from the product space  $X \times Y$  onto  $Y$  is closed for any paracompact Hausdorff space  $Y$  such that each point of  $Y$  has a neighborhood basis of power  $\leq m$ .*

*Proof.* As the "only if" part has been shown in our previous note,<sup>2)</sup> we need only prove the "if" part. If we suppose that  $X$  is not  $m$ -compact, then there exists a collection of closed subsets  $\mathfrak{F} = \{F_\lambda \mid \lambda \in \Lambda\}$  with the finite intersection property such that

- (1)  $|\Lambda| \leq m$  where  $|\Lambda|$  denotes the power of  $\Lambda$ .
- (2)  $\bigcap_{\lambda \in \Lambda} F_\lambda = \phi$ .

Moreover, by adding to  $\mathfrak{F}$  all the intersections of finitely many sets of  $\mathfrak{F}$ , we can assume that  $\mathfrak{F}$  satisfies the following condition (3), because  $|\Lambda|$  does not exceed  $m$ .

- (3)  $F_\lambda \cap F_\mu \in \mathfrak{F}$  for any two sets  $F_\lambda, F_\mu$  of  $\mathfrak{F}$ .

We define the partial order in such a way that  $\lambda \geq \mu$  if and only if  $F_\lambda \subset F_\mu$ . Then  $\Lambda$  is a directed set by the condition (3).

Let  $Y$  denote the set of different elements  $\{y_\lambda \mid \lambda \in \Lambda\} \cup y_\infty$ , where  $\infty \neq \lambda$  for every  $\lambda \in \Lambda$ . We next define the topology of  $Y$  such that

- (4) the neighborhood basis of each point  $y_\lambda$  is the single point set  $\{y_\lambda\}$ ,
- (5) the neighborhood basis of the point  $y_\infty$  is the family of sets  $U_\lambda(y_\infty) = \{y_\mu \mid \mu \geq \lambda\} \cup y_\infty$ .

Then, since  $\Lambda$  is a directed set,  $Y$  is a topological space. It is evident that each point of  $Y$  has a neighborhood basis of power  $\leq m$ . We next prove that  $Y$  is a Hausdorff space. Since  $\{y_\lambda\} \cap \{y_\mu\} = \phi$

1) K. Nagami communicated to me this interesting theorem in his kind letter of August 8, 1961.

2) S. Hanai: Inverse images of closed mappings. I, Proc. Japan Acad., **37**, 298-301 (1961).