

80. On the Regularity of Domains for Parabolic Equations

By Haruo MURAKAMI

Kobe University

(Comm. by K. KUNUGI, M.J.A., June 12, 1958)

Let G be a domain with the boundary Γ in the m -dimensional Euclidean space R^m . Let $D = G \times (0, \infty)$ and $B = \Gamma \times [0, \infty)$. W. Fulks pointed out in [1] by constructing barriers at every boundary point of G that if D is regular for the heat equation then G is regular for Laplace's equation. In this note we shall show also by constructing barriers of the parabolic equation at every point of $G \cup B$ that the converse of the result above by W. Fulks is true.

Consider the equation

$$(E) \quad \square u = f(x, t, u) \quad 1)$$

where $f(x, t, u)$ is continuous on $D \times (0, \infty)$ and quasi-bounded with respect to u .

As in [2, p. 623], we say that $w(x, t)$ is a barrier of (E) at a boundary point $(x^0, t^0) \in G \cup B$ with respect to a bounded function $\beta(x, t)$ defined on $G \cup B$ if $w(x, t)$ satisfies:

- (i) $w(x, t)$ is continuous on \bar{D} ,
- (ii) $w(x, t) > 0 \quad (x, t) \in \bar{D}, \quad (x, t) \neq (x^0, t^0)$,
- (iii) $w(x, t) \rightarrow 0 \quad (x, t) \rightarrow (x^0, t^0), \quad (x, t) \in \bar{D}$,
- (iv) $\square w(x, t) \leq -M$, where

$$M = \sup \{ |f(x, t, \bar{\beta}(x^0, t^0))|, |f(x, t, \underline{\beta}(x^0, t^0))|; (x, t) \in \bar{D} \}.$$

It is known²⁾ that if every point of $G \cup B$ has barriers then D is regular for (E), i.e. the first boundary value problem of (E) is always solvable for any continuous data on $G \cup B$.

Now we shall construct the barrier $w(x, t)$ satisfying the conditions (i), (ii), (iii) and (iv) under the assumption that G is regular for Laplace's equation.

In case that $(x^0, t^0) \in G$, it is easy to see that the function $w(x, t) = \sum_{i=1}^m (x_i - x_i^0)^2 + (2m + M)(t - t^0)$ is a barrier at (x^0, t^0) . In case that $(x^0, t^0) \in B$ with $t^0 > 0$, let $\varphi(x, t) = \sum_{i=1}^m (x_i - x_i^0)^2 + (t - t^0)^2$. Then we have $\varphi(x, t) \geq 0$ and

$$\begin{aligned} \square \varphi(x, t) &= \sum_{i=1}^m \frac{\partial^2 \varphi(x, t)}{\partial x_i^2} - \frac{\partial \varphi(x, t)}{\partial t} \\ &= 2(m - (t - t^0)). \end{aligned}$$

1) \square and \triangle below are respectively the generalized heat and Laplacian operators. For the definitions, see [2, p. 627], where \square is denoted by \square . See also [3, p. 349].

2) [2, pp. 624-626].