

## 29. Total Orderings on a Semilattice<sup>1)</sup>

By Naoki KIMURA

Tokyo Institute of Technology and Tulane University

(Comm. by K. SHODA, M.J.A., March 12, 1958)

A semilattice  $S$  is called *orderable* if there exists an ordering  $\leq$  such that (1)  $a < b$  implies  $ac \leq bc$  and (2)  $a \leq b$  or  $b \leq a$  for any  $a, b \in S$ . Such an ordering is called *permissible* on  $S$ . The main purpose of this note is to present a necessary and sufficient condition for a semilattice to be orderable.<sup>2)</sup>

**THEOREM 1.** *A semilattice is orderable if and only if it does not contain any subsemilattice consisting of four elements  $a, b, c, d$  satisfying either (i)  $ab=b, ac=c, bc=d$  or (ii)  $ab=ac=bc=d$ .*

**COROLLARY.** *Any chain, in the sense that  $ab=a$  or  $b$  for all  $a, b$ , is orderable.*

An element  $a$  of a semilattice is called *maximal* if  $ax=a$  implies  $x=a$ .

**THEOREM 2.** *Let  $S$  be an orderable semilattice. Let  $T$  be the complement of the set of all maximal elements in  $S$ . Then there exists a one to one correspondence between the set of all permissible orderings on  $S$  and the set of all subsets of  $T$ .*

**COROLLARY.** *Let  $N(n)$  be the number of all non-isomorphic orderable semilattices consisting of  $n$  elements. Then it satisfies the following formula:*

$$N(n+1) = \sum_{\substack{0 \leq p < q \leq n \\ p+q=n}} N(p)N(q) + \begin{cases} \frac{1}{2}N(n/2)(N(n/2)+1) & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Further,  $N(n)$  is equal to the coefficient of  $x^n$  in the expansion of  $f(x)$  defined by

$$f(x^2) + x(f(x))^2 - 2f(x) + 2 = 0.$$

We shall close this note by listing here two examples as application.

**EXAMPLE 1.** Let  $S$  be a chain. Let  $(A, B)$  be a partition of  $S$ , in the sense that  $S=A \cup B$ ,  $A \cap B = \square$ , the empty set. Then the ordering defined by

$$x \leq y \quad \text{if and only if} \quad \begin{cases} x, y \in A, & xy = x \\ \text{or } x \in A, & y \in B \\ \text{or } x, y \in B, & xy = y, \end{cases}$$

gives a permissible ordering on  $S$ .

Conversely, any permissible ordering on  $S$  can be obtained by a

1) This work was partially supported by the National Science Foundation, U. S. A.

2) This is an abstract of the paper which will appear elsewhere.