

155. A Characterisation of Regular Semi-group

By Kiyoshi ISÉKI

(Comm. by K. KUNUGI, M.J.A., Nov. 12, 1956)

The concept of regular ring was introduced by J. v. Neumann [3]. Recently, L. Kovács [1] has given an interesting characterisation of regular ring. On the other hand, some writers studied regular semi-groups. In this short Note, we shall give a characterisation for regular semi-group which is similar to Theorem 1 of L. Kovács [1].

Following W. D. Munn and R. Penrose [2], we define a regular semi-group. A semi-group is said to be *regular* if for any given element a of S there is at least one element x of S such that $axa=a$. A non empty subset L of S is said to be a *left ideal* if $SL \subset L$. Similarly we define *right ideals*. Then we have the following

Theorem 1. Any semi-group S is regular if and only if

$$AB = A \cap B$$

for every right ideal A and every left ideal B of S .

Proof. Let S be a regular semi-group, and let $a \in A \cap B$, then there is an element x such that $axa=a$. Since B is a left ideal, $xa \in B$. Therefore $a = a(xa) \in AB$. This shows $AB \supset A \cap B$. Clearly $AB \subset A \cap B$. Hence $AB = A \cap B$.

To prove the converse, let a be an element of S . Then $\{ax \mid x \in S\} \cup a$ is the right ideal (a) of S generated by a . By the hypothesis,

$$(a) = (a) \cap S = (a)S = aS.$$

Therefore, we have $a \in aS$. Similarly $a \in Sa$. Hence

$$a \in aR \cap Ra = aR^2a,$$

and there is an element x such that $a = axa$.

Now, let us suppose that a given regular semi-group S is *commutative*, then, by Theorem 1, any ideal A in S is idempotent, i.e. $A^2 = A$. Conversely, suppose that every ideal in a commutative semi-group S is idempotent. If A and B are ideals in S , then we have $A \cap B = (A \cap B)^2 = (A \cap B)(A \cap B) \subset AB$. On the other hand, $A \cap B \supset AB$. Hence $A \cap B = AB$. By Theorem 1, S is regular, therefore we have the following

Theorem 2. A commutative semi-group is regular if and only if every ideal is idempotent.

From Theorem 2, it is easily seen that there is no non-zero nilpotent element in a commutative regular semi-group with 0.

Corollary. Any commutative regular semi-group with 0 does not