

### 109. A Remark on Theorems of Stone and Bochner.

By Masahiro NAKAMURA and Hisaharu UMEGAKI.

Osaka Normal College and Mathematical Institute, Kyusyu University.

(Comm. by K. KUNUGI, M.J.A., Nov. 12, 1951.)

Recent progress of the theory of topological groups generalizes many theorems of real analysis into locally compact abelian groups. Among them, Raikov-Ambrose's extension of the well-known theorem of Stone reads as follows: *Let  $g^*$  is a continuous unitary representation of a locally compact abelian group  $G$  on a Hilbert space  $H$ . Then there exists a resolution of the identity  $de(\chi)$  which is defined for all Borel sets of the character group  $G^*$  such that*

$$(1) \quad g^* = \int \chi(g) de(\chi).$$

Combining this and the well-known theorem of Gelfand-Raikov which gives the representation of positive definite functions on  $G$ , we can deduce Bochner-Weil-Raikov's representation of positive definite functions putting  $d\mu(\chi) = d(\xi e(\chi), \xi)$ ; that is, *each positive definite function  $\varphi(g)$  allows a representation*

$$(2) \quad \varphi(g) = \int \chi(g) d\mu(\chi),$$

where  $d\mu(\chi)$  means a regular Borel measure on  $G^*$ . Therefore if we can prove Stone's theorem directly, then we can prove Bochner's theorem as in the above.

In this note, we shall do this utilizing recent published theorem of N. Dunford [1], which states that *a commutative  $C^*$ -algebra  $A$  (in the sense of I. E. Segal [2]; i.e., a uniformly closed operator algebra on a Hilbert space  $H$ ) admits the integral representation*

$$(3) \quad x = \int_{\Omega} x(\omega) de(\omega),$$

where  $\Omega$  is the compact space of all maximal ideals of  $A$  (with the Stone topology),  $x(\omega)$  is the representation of  $x$  on  $\Omega$  and  $de(\omega)$  is the resolution of the identity defined for all Borel sets of  $\Omega$ . Although Dunford proved the theorem for compact space, it is not difficult to generalize for locally compact case deleting the identity (we will use this generalized form in the below).

Let  $R(G)$  be the operator group algebra of  $G$  in the sense of I. E. Segal [2], i.e., the uniform closure of the "regular" representation of  $L_1(G)$ . Furthermore suppose  $A$  be the uniform closure of the operators of the form :