

## 85. A Note on Hausdorff Spaces with the Star-finite Property. III

By Keiô NAGAMI

Ehime University, Matsuyama

(Comm. by K. KUNUGI, M.J.A., July 12, 1961)

We shall prove in this note, by a very simple argument, that an arbitrary non-empty (separable) metric space  $R$  is the image of a 0-dimensional (separable) metric space, under the open continuous mapping. At the first sight this is an odd fact, in view of Yu. Rozanskaya's theorem [3] which asserts that there does not exist an open continuous mapping of an  $m$ -dimensional Euclidean cube  $R_m$  onto an  $n$ -dimensional Euclidean cube  $R_n$  with  $m < n$ .

**Theorem 1.** *A topological  $T_1$ -space  $R$  is always the image of a completely regular space  $A$  with  $\text{ind } A = 0$  under the open continuous mapping  $f$  such that  $f^{-1}(x)$  is compact for every point  $x$  of  $R$ .*

*Proof.* Let  $\{\mathfrak{U}_\lambda = \{U_\alpha; \alpha \in A_\lambda\}; \lambda \in \Lambda\}$  be a family of all finite open coverings of  $R$ . Let  $A$  be the aggregate of points  $a = (\alpha_i; \lambda \in \Lambda)$  of the product space  $\Pi \{A_\lambda; \lambda \in \Lambda\}$ , where  $A_\lambda$  are topological spaces with the discrete topology, such that  $\bigcap \{U_\alpha; \lambda \in \Lambda\} \neq \emptyset$ . Let  $f(a) = \bigcap \{U_{\pi_\lambda(a)}; \lambda \in \Lambda\}$ , where  $\pi_\lambda: A \rightarrow A_\lambda$ ,  $\lambda \in \Lambda$ , are the projections. Then  $f$  is a mapping of  $A$  onto  $R$ . Since for any  $\lambda \in \Lambda$  and any  $\alpha \in A_\lambda$  we have  $f(\pi_\lambda^{-1}(\alpha)) = U_\alpha$ ,  $f$  is an open continuous mapping. Let  $x$  be an arbitrary point of  $R$  and  $B_\lambda = \{a; x \in U_\alpha \in \mathfrak{U}_\lambda\}$ ,  $\lambda \in \Lambda$ . Then  $f^{-1}(x) = \bigcap B_\lambda$  and hence it is compact. It is almost evident that  $A$  is a completely regular space with  $\text{ind } A = 0$ . Thus the theorem is proved.

**Theorem 2.** *A non-empty metric space  $R$  is always the image of a metric space  $A$  with  $\text{dim } A = 0$ , under the open continuous mapping  $f$  such that  $f^{-1}(x)$  is compact for every point  $x$  of  $R$ .*

*Proof.* Since a metric space is always paracompact by A. H. Stone [4, Corollary 1], there exists a sequence  $\mathfrak{U}_i = \{U_\alpha; \alpha \in A_i\}$ ,  $i = 1, 2, \dots$ , of locally finite open coverings of  $R$  such that the diameter of each element of  $\mathfrak{U}_i$  is less than  $1/i$ . Let  $A$  be the aggregate of points  $a = (\alpha_i; i = 1, 2, \dots)$  of the product space  $\Pi \{A_i; i = 1, 2, \dots\}$ , where  $A_i$  are topological spaces with the discrete topology, such that  $\bigcap \{U_\alpha; i = 1, 2, \dots\} \neq \emptyset$ . Let  $f(a) = \bigcap \{U_{\pi_i(a)}; i = 1, 2, \dots\}$ , where  $\pi_i: A \rightarrow A_i$ ,  $i = 1, 2, \dots$ , are the projections. Then by the same argument as in the proof of Theorem 1  $f$  becomes an open continuous mapping of  $A$  onto  $R$  such that  $f^{-1}(x)$  is compact for every point  $x$  of  $R$ . Moreover  $A$  is a metric space with  $\text{dim } A = 0$  by Katětov [1, Theorem 3.7] or Morita [2, Theorem 10.2]. Thus the theorem is proved.