

## 142. A Note on the Number of Irreducible Characters in a $p$ -Block with Normal Defect Group

By Masafumi MURAI

(Communicated by Shokichi IYANAGA, M. J. A., Dec. 12, 1983)

1. Let  $G$  be a finite group and  $p$  be a prime. Let  $B$  be a  $p$ -block of  $G$  with defect group  $D$ . We denote by  $k(B)$  the number of ordinary irreducible characters in  $B$ . R. Brauer [1] conjectured

$$(K): k(B) \leq |D|.$$

In [5] it is shown that (K) is true if  $G$  is  $p$ -solvable and if  $p$  is sufficiently large compared with the sectional rank of  $D$ .

The purpose of this note is to prove the following

**Theorem.** *For any positive integer  $n$ , there exists a constant  $b_n$  depending only on  $n$  such that the following statement is true: Let  $B$  be a  $p$ -block of a group  $G$  with normal defect group  $D$ . Assume that the sectional rank of  $D$  equals  $n$ . Then, if  $p$  is larger than  $b_n$ , we have  $k(B) \leq |D|$ .*

2. Let  $B$  be a  $p$ -block of a group  $G$  with defect group  $D$ , which is normal in  $G$ . Let  $b$  be a  $p$ -block of  $DC_a(D)$  covered by  $B$  and  $T_b$  be the inertia group in  $G$  of the block  $b$ . Then  $[T_b : DC_a(D)]$  is prime to  $p$ . Let  $B'$  be the unique block of  $T_b$  that covers  $b$ . Then  $D$  is the defect group of  $B'$  and  $k(B') = k(B)$ . In order to prove that (K) is true for  $B$  we may assume that  $G = T_b$ ,  $B = B'$ . Then  $G/C_a(D)$  contains the normal  $p$ -Sylow group  $DC_a(D)/C_a(D)$ , so that  $G/C_a(D)$  has a  $p$ -complement  $L/C_a(D)$ . Set  $\bar{L} = L/C_a(D)$ . Form the semi-direct product  $H = \bar{L}D$  with the natural (faithful) action of  $\bar{L}$  on  $D$ . Theorem follows immediately from the result in [5] mentioned above and the following

**Proposition.** *Let the notation be as above. We have  $k(B) \leq \text{cl}(H)$ . Here  $\text{cl}(X)$  denotes the number of conjugacy classes of  $X$  for a group  $X$ .*

*Proof.* Let  $\theta$  be the canonical character of  $b$ . For every irreducible character  $\chi$  of  $D$ , define the class function  $\tilde{\chi}$  on  $DC_a(D)$  as follows:

$$\tilde{\chi}(z) = \begin{cases} \chi(x)\theta(y) & \text{if } z = xy \text{ with } x \in D, y \in C_a(D) \\ 0 & \text{otherwise,} \end{cases}$$

where  $x$  and  $y$  denote the  $p$ -part and  $p'$ -part of  $z \in DC_a(D)$ , respectively. Then the map  $\sim$  is a bijection from the set of irreducible characters of  $D$  onto the set of irreducible characters in  $b$  (see [2], (V. 4.7)). Let  $\{\chi_i\}$  be a complete set of representatives of  $\bar{L}$ -conjugate