

97. Spectral Properties of Random Media

By Shin OZAWA

Department of Mathematics, Osaka University

(Communicated by Kôzaku YOSIDA, M. J. A., Dec. 12, 1984)

This note is a continuation of our previous papers Ozawa [2], [3]. We consider a bounded domain Ω in \mathbf{R}^3 with smooth boundary Γ . We put $B(\varepsilon; w) = \{x \in \mathbf{R}^3; |x - w| < \varepsilon\}$. Fix $\beta \geq 1$. Let $0 < \mu_1(\varepsilon; w(m)) \leq \mu_2(\varepsilon; w(m)) \leq \dots$ be the eigenvalues of $-\Delta (= -\text{div grad})$ in $\Omega_{\varepsilon, w(m)} = \Omega \setminus \bigcup_{i=1}^{\tilde{m}} B(\varepsilon; w_i^{(m)})$ under the Dirichlet condition on its boundary. Here \tilde{m} denotes the largest integer which does not exceed m^β , and $w(m)$ denotes the set of \tilde{m} -points $\{w_i^{(m)}\}_{i=1}^{\tilde{m}} \in \Omega^{\tilde{m}}$. Let $V(x) > 0$ be C^1 -class function on $\bar{\Omega}$ satisfying

$$\int_{\Omega} V(x) dx = 1.$$

We consider Ω as the probability space with the probability density $V(x) dx$. Let $\Omega^{\tilde{m}} = \prod_{i=1}^{\tilde{m}} \Omega$ be the probability space with the product measure.

In this note we give the following:

Theorem 1. Fix $\beta \in [1, 3)$. Assume that $V(x) > 0$. Fix $\alpha > 0$ and k . Then, there exists a constant $\delta(\beta) > 0$ independent of m such that

$$\lim_{m \rightarrow \infty} P(w(m) \in \Omega^{\tilde{m}}; m^{\delta' - (\beta - 1)} |\mu_k(\alpha/m; w(m)) - \mu_{k,m}^V| < \varepsilon) = 1$$

holds for any $\varepsilon > 0$ and $\delta' \in [0, \delta(\beta))$. Here $\mu_{k,m}^V$ denotes the k -th eigenvalue of $-\Delta + 4\pi\alpha m^{\beta-1} V(x)$ in Ω under the Dirichlet condition on Γ .

We give a short sketch of our proof of Theorem 1. In the following we use notations and terminologies in [2]. We have the following:

Lemma 1. Fix $\beta \in [1, 3)$. Suppose that $u_m \in C^\infty(\omega)$ satisfies

$$\begin{aligned} (-\Delta + m')u_m(x) &= 0, & x \in \omega \\ u_m(x) &= 0, & x \in \partial\omega \cap \Gamma \end{aligned}$$

and

$$\max \{|u_m(x)|; x \in \partial B_r \cap \partial\omega\} = M_r(m),$$

$r = 1, \dots, \tilde{m}$. If $\partial B_r \cap \partial\omega = \phi$, then we put $M_r(m) = 0$. Under the above assumption, there exists a constant C independent of m such that

$$\begin{aligned} \|u\|_{L^2(\omega)} &\leq C(\alpha/m)(m')^{-1/4} \left(\sum_{r=1}^{\tilde{m}} M_r(m)^2 \right)^{1/2} \\ &\quad + C(\alpha/m)(m')^{-1/4} \left(\sum_{\substack{r,s=1 \\ r \neq s}}^{\tilde{m}} \exp(- (m')^{1/2} |w_r - w_s|) M_r(m) M_s(m) \right)^{1/2} \end{aligned}$$

holds.

By using Lemma 1 we get the following;