

16. Modular Pairs in the Lattice of Projections of a von Neumann Algebra

By Shūichirō MAEDA

Department of Mathematics, Ehime University

(Communicated by Kōsaku YOSIDA, M. J. A., Feb. 12, 1986)

A pair (a, b) of elements of a lattice is called modular (resp. dual-modular), denoted by $(a, b)M$ (resp. $(a, b)M^*$), if

$$(c \vee a) \wedge b = c \vee (a \wedge b) \text{ for every } c \leq b \\ \text{(resp. } (c \wedge a) \vee b = c \wedge (a \vee b) \text{ for every } c \geq b)$$

(see [3], (1.1)). If A is a von Neumann algebra, then the set $P(A)$ of all projections of A forms an orthomodular lattice ([3], (37.13) and (37.15)). An algebraic characterization of modular pairs of projections is given in Section 38 of [3]. In this paper, we shall give a norm characterization of modular pairs in $P(A)$, by using the result of Bures [1].

Lemma 1. *Let a, b be elements of an orthomodular lattice, and we put $a_0 = a - a \wedge b$, $b_0 = b - a \wedge b$ (see [3], (36.5)). Then,*

$$(a, b)M \iff (a_0, b_0)M \iff (a, b_0)M.$$

Proof. Assume $(a_0, b_0)M$. Since $a_0 \leq (a \wedge b)^\perp$, we have $a_0 C a \wedge b$ by [3], (36.3). Similarly, $b_0 C a \wedge b$. Hence, it follows from [3], (36.11) that $(a_0 \vee (a \wedge b), b_0 \vee (a \wedge b))M$, which implies $(a, b)M$.

Next, if we assume $(a, b)M$, then since $a C (a \wedge b)^\perp$, $b C (a \wedge b)^\perp$ and $b \wedge (a \wedge b)^\perp = b_0$, we have $(a, b_0)M$ by [3], (36.11). Finally, if we assume $(a, b_0)M$, then since $a \wedge b_0 = (a \wedge b) \wedge b_0 \leq (a \wedge b) \wedge (a \wedge b)^\perp = 0$, we have $(a_0, b_0)M$ by [3], (1.5.3).

In [1], two elements a and b of a lattice are said to be modularly separated, if $a \wedge b = 0$ and $(a', b')M^*$, $(b', a')M^*$ for all $a' \leq a$ and $b' \leq b$.

Lemma 2. *Let A be a von Neumann algebra and let $e, f \in P(A)$.*

(i) $(e, f)M \iff (f, e)M \iff (e, f)M^* \iff (f, e)M^*$.

(ii) *e and f are modularly separated if and only if $e \wedge f = 0$ and $(e, f)M$.*

Proof. Since A is a Baer *-ring satisfying the condition (SR) ([3], (37.15)), (i) follows from [3], (29.8) and (37.14). The "only if" part of (ii) follows from (i). Conversely, if $e \wedge f = 0$ and $(e, f)M$ then $(e', f')M$ for all $e' \leq e$, $f' \leq f$ by [3], (1.5.3). Hence, e and f are modularly separated by (i).

We remark that Theorem 6 and Corollary of Theorem 7 in [1] immediately follow from our Lemma 2 (ii) and [3], (1.6).

Theorem 3. *Let A be a von Neumann algebra and let $e, f \in P(A)$. (e, f) is a modular pair in $P(A)$ if and only if there exist a finite element $e_1 \in P(A)$ with $e_1 \leq e - e \wedge f$ and an orthogonal sequence of central projections*