

### 46. The Flat Holomorphic Conformal Structure on the Horrocks-Mumford Orbifold

By Takeshi SATO

Department of Mathematics, Faculty of Science, University of Tokyo

(Communicated by Kunihiko KODAIRA, M. J. A., May 13, 1991)

We construct the explicit flat holomorphic conformal structure on an orbifold. We often abbreviate ‘Horrocks-Mumford’ to ‘HM’, and ‘holomorphic conformal structure’ to ‘HCS’.  $P_n(C)$  denotes the  $n$ -dimensional complex projective space.

In the paper [2], Horrocks and Mumford constructed a holomorphic vector bundle  $\mathcal{F}_{HM}$  of rank two on the  $P_4(C)$ . The space  $\Gamma\mathcal{F}_{HM}$  of its sections is four-dimensional. If the zero set  $X_s$  of a section  $s \in \Gamma\mathcal{F}_{HM}$  is a smooth surface,  $X_s$  is an abelian surface with (1.5)-polarization. In fact, they proved that  $P_3(C) = P(\Gamma\mathcal{F}_{HM})$  is birationally equivalent to the moduli space  $\mathcal{A}_{1,5}$  of the abelian surfaces with (1, 5)-polarization and level-5-structure. (See [2] [4].) We call this projective space *the HM-orbifold*.

While the moduli space  $\mathcal{A}_{1,5}$  is realized as a quotient space  $\mathcal{H}_2/\Gamma_{1,5}$  of the Siegel upper space  $\mathcal{H}_2$  of degree two. Here  $\Gamma_{1,5}$  is a certain discrete subgroup of  $Sp(4, \mathbf{R})$ . (See [4].)  $\mathcal{H}_2$  is embedded in a non-degenerate hyperquadrics  $\{[z_0 : z_1 : z_2 : z_3 : z_4] \in P_4(C) ; \sum_{0 \leq i, j \leq 4} a_{ij} z_i z_j = 0\}$ . The holomorphic tensor field  $\phi = \sum_{0 \leq i, j \leq 4} a_{ij} dz_i dz_j$  on  $\mathcal{H}_2$  is conformally flat and its conformally class is invariant under the automorphisms of  $\mathcal{H}_2$ . Therefore  $\phi$  induces a tensor  $\varphi$  on the HM-orbifold which is called *the flat HCS*. Applying a higher dimensional version of Kobayashi and Naruki’s theory in [3], we can calculate the flat HCS.

**Theorem 1.** *Let  $p$  be the projection  $C^4 \setminus \{0\} \rightarrow P_3(C)$ . The pullback of the flat HCS  $\varphi$  is given by in homogeneous coordinates*

$$p^*\varphi = \sum_{0 \leq i, j \leq 3} g_{ij} dx_i dx_j$$

where

$$\begin{aligned} g_{00} &= 2(-x_0^2 x_1 x_2 - x_0^2 x_3^2 + x_0 x_1^3 + 2x_0 x_2^2 x_3 + 2x_1^2 x_2 x_3 - 3x_1 x_3^3) \\ g_{01} &= x_0^3 x_2 - 2x_0^2 x_1^2 - 7x_0 x_1 x_2 x_3 + 4x_0 x_3^3 + x_1^3 x_3 + 4x_1 x_2^3 - 5x_2^2 x_3^2 \\ g_{02} &= x_0^3 x_1 - x_0^2 x_2 x_3 - x_0 x_1^2 x_3 - 4x_1^2 x_2^2 + 5x_1 x_2 x_3^2 \\ g_{03} &= 2x_0^3 x_3 - 3x_0^2 x_2^2 + 4x_0 x_1^2 x_2 + 2x_0 x_1 x_3^2 - x_1^4 \\ g_{11} &= 2(x_0^3 x_1 + 2x_0^2 x_2 x_3 - x_0 x_1^2 x_3 - 3x_0 x_2^3 - x_1^2 x_2^2 + 2x_1 x_2 x_3^2) \\ g_{12} &= -x_0^4 + 4x_0^2 x_1 x_3 + 2x_0 x_1 x_2^2 + 2x_1^3 x_2 - 3x_1^2 x_3^2 \\ g_{13} &= -x_0^2 x_1 x_2 - 4x_0^2 x_3^2 + x_0 x_1^3 + 5x_0 x_2^2 x_3 - x_1^2 x_2 x_3 \\ g_{22} &= 2(-x_0^3 x_3 + x_0 x_1^2 x_2 + 5x_0 x_1 x_3^2 - x_1^4) \\ g_{23} &= 3(x_0^3 x_2 - x_0^2 x_1^2 - 5x_0 x_1 x_2 x_3 + x_1^3 x_3) \\ g_{33} &= 2(-x_0^4 + x_0^2 x_1 x_3 + 5x_0 x_1 x_2^2 - x_1^3 x_2) \\ g_{10} &= g_{01}, g_{20} = g_{02}, g_{30} = g_{03}, g_{21} = g_{12}, g_{31} = g_{13}, g_{32} = g_{23}. \end{aligned}$$