

67. Absolutely Continuous Spectra of Relativistic Schrödinger Operators with Magnetic Vector Potentials

By Tomio UMEDA

Department of Mathematics, Himeji Institute of Technology
(Communicated by Kiyosi ITÔ, M. J. A., Nov. 14, 1994)

Key words: Spectral theory; scattering theory; relativistic spinless particles; magnetic vector potential.

§1. Introduction. A spinless particle in an electromagnetic field is described by a relativistic Schrödinger operator

$$H = h^w(x, D) + V(x),$$

where

$$h^w(x, D)u(x) = \int \int e^{i(x-y) \cdot \xi} h\left(\frac{x+y}{2}, \xi\right)u(y) (2\pi)^{-n} dy d\xi,$$

$$h(x, \xi) = \sqrt{|\xi - a(x)|^2 + m^2},$$

$$(x, \xi \in \mathbf{R}^n, m > 0).$$

Although many efforts have been made to understand the nature of the operator H , there are few works in which spectral properties of H are investigated (cf. [2]-[7]). Indeed, Nagase and Umeda [8] is the only work locating the spectrum of the operator H as far as we know. In [8] they showed that $\sigma_{\text{ess}}(H) = [m, \infty)$ under the following assumptions (precisely speaking, their assumptions on $a(x)$ are weaker than (I) and (II) below):

- (I) For any multi-index α , $\partial_x^\alpha a(x) \rightarrow 0$ as $|x| \rightarrow \infty$.
- (II) $V(-\Delta + 1)^{-1/2}$ is a compact operator in $L^2(\mathbf{R}^n)$.

From the view point of spectral and scattering theory, it is natural to ask whether the absolutely continuous spectrum of H coincides with the interval $[m, \infty)$, and whether the singular continuous spectrum is empty. The aim of this note is to make a remark that the Enss theory can give answers to these questions. Finally, we mention here that our assumption on $a(x)$ is stronger than (I) and (II) above for technical reason.

§2. Results. Throughout this section, ε denotes a positive constant and $\langle x \rangle = \sqrt{1 + |x|^2}$. For the vector and scalar potentials $a(x) = (a_1(x), \dots, a_n(x))$ and $V(x)$, we make the following assumptions respectively:

- (A) Each $a_j(x)$ is a C^∞ -function such that $|\partial^\alpha a_j(x)| \leq C_\alpha \langle x \rangle^{-1-\varepsilon}$ for any α .
- (V) $V(x)$ is a real-valued measurable function such that $|V(x)| \leq C \langle x \rangle^{-1-\varepsilon}$.

It is known [8, section 2] that under assumptions (A) and (V) the operator $h^w(x, D) + V(x)$ restricted on $C_0^\infty(\mathbf{R}^n)$ is essentially self-adjoint in $L^2(\mathbf{R}^n)$. Its self-adjoint realization will be denoted by H again. It is also known [8, section 2] that $\text{Dom}(H) = H^1(\mathbf{R}^n)$, the Sobolev space of order 1. Our result is

Theorem. *Let (A) and (V) be satisfied. Then*