

A Note on the Capitulation in \mathbf{Z}_p -extensions

By Manabu OZAKI

Department of Mathematics, School of Science and Engineering, Waseda University

(Communicated by Shokichi IYANAGA, M. J. A., Nov. 13, 1995)

1. Introduction. Let K be a number field, namely, a finite extension field over the field of rational numbers. Let p be a prime number, and K_∞/K a \mathbf{Z}_p -extension with Galois group Γ . Denote by K_n the n -th layer of K_∞/K . We write A_n for the p -Sylow subgroup of the ideal class group of K_n , and $j_{n,m}$ for the map from A_n to A_m induced from the inclusion $K_n \subseteq K_m$ for $m \geq n \geq 0$. Let $A_\infty = \varinjlim A_n$, where the inductive limit is taken with respect to $j_{n,m}$, and let $j_{n,\infty}$ be the natural map from A_n to A_∞ . We denote by $\lambda(K_\infty/K)$, $\mu(K_\infty/K)$ the Iwasawa λ, μ invariant, respectively, of K_∞/K .

In the present paper, we shall prove the following:

Theorem. *Let notations be as above. If we assume that all primes of K which ramify in K_∞ are totally ramified in K_∞ , then*

$$\lambda(K_\infty/K) = \mu(K_\infty/K) = 0 \Leftrightarrow \text{Ker}(N_{1,0}: A_1 \rightarrow A_0) \subseteq \text{Ker}(j_{1,\infty}: A_1 \rightarrow A_\infty),$$

where $N_{1,0}$ stands for the norm map from A_1 to A_0 .

This theorem is in analogy to the following theorem due to Greenberg [1, Theorem 1], though our proof is based on a method different from [1].

Theorem (Greenberg). *Let K be a totally real number field, and K_∞/K a \mathbf{Z}_p -extension. If we assume that there is only one prime in K above p , which is totally ramified in K_∞ , then*

$$\lambda(K_\infty/K) = \mu(K_\infty/K) = 0 \Leftrightarrow \text{Ker}(j_{0,\infty}: A_0 \rightarrow A_\infty) = A_0.$$

2. Proof of Theorem. We fix a topological generator γ of Γ . Put $\Lambda = \mathbf{Z}_p[[\Gamma]]$ and $\nu_{n,m} = \frac{\gamma^{p^m} - 1}{\gamma^{p^n} - 1} \in \Lambda$ for $m \geq n \geq 0$. Let L_∞/K_∞ be the maximal unramified pro- p abelian extension, and let X be its Galois group. Put $Y = \text{Gal}(L_\infty/K_\infty L_0)$, where L_0 is the Hilbert p -class field of K . Then X is a finitely generated torsion Λ -module, and the Artin map induces the isomorphism $A_n \cong X/\nu_{0,n}Y$ for all $n \geq 0$ (cf. [2, Theorem 6]). So we will identify $X/\nu_{0,n}Y$ with A_n .

We first prove the following:

Proposition. $\text{Ker}(j_{n,\infty}: A_n \rightarrow A_\infty) = \text{Im}(X_{finite})$

$\rightarrow X/\nu_{0,n}Y$) for all $n \geq 0$, where X_{finite} denotes the maximal finite Λ -submodule of X .

Proof. Since X_{finite} is finite, we see that $\nu_{n,m}X_{finite} = 0$ for some $m \geq n$. Observing the following commutative diagram (cf. [2, Theorem 7]):

$$(1) \quad \begin{array}{ccc} A_m & \xrightarrow{\sim} & X/\nu_{0,m}Y \\ j_{n,m} \uparrow & & \uparrow \nu_{n,m} \\ A_n & \xrightarrow{\sim} & X/\nu_{0,n}Y, \end{array}$$

we have

$$\text{Im}(X_{finite} \rightarrow X/\nu_{0,n}Y) \subseteq \text{Ker}(j_{n,\infty}: A_n \rightarrow A_\infty).$$

Conversely, let $x \text{ mod. } \nu_{0,n}Y \in X/\nu_{0,n}Y$ be any element in $\text{Ker}(j_{n,\infty}: A_n \rightarrow A_\infty)$. It follows from (1) that $\nu_{n,m}x \in \nu_{0,m}Y$ for some $m \geq n$. Hence $\nu_{n,m}x = \nu_{0,m}y$ for some $y \in Y$, that is

$$(2) \quad \nu_{n,m}(x - \nu_{0,n}y) = 0.$$

X/X_{finite} is embedded in the Λ -module $\bigoplus_{i=1}^r \Lambda/f_i^e \Lambda$ with finite cokernel, where $f_i \in \Lambda$ is a prime element. Since $X/\nu_{n,m}X$ is finite, $\prod_{i=1}^r f_i^{e_i}$ is prime to $\nu_{n,m}$. So we see that the multiplication-by- $\nu_{n,m}$ map $\nu_{n,m}: X/X_{finite} \rightarrow X/X_{finite}$ is injective. Therefore we have $x - \nu_{0,n}y \in X_{finite}$ from (2). So we obtain $x \text{ mod. } \nu_{0,n}Y \in \text{Im}(X_{finite} \rightarrow X/\nu_{0,n}Y)$. \square

Corollary. $\text{Ker}(j_{n,\infty}: A_n \rightarrow A_\infty) \neq 0$ for some $n \geq 0 \Leftrightarrow X_{finite} \neq 0$

Proof. (\Rightarrow) part is obvious by Proposition. We assume that $X_{finite} \neq 0$. It follows from $\bigcap_{n \geq 0} \nu_{0,n}Y = 0$ that $X_{finite} \not\subseteq \nu_{0,n}Y$ for some $n \geq 0$. Therefore we find that $\text{Ker}(j_{n,\infty}: A_n \rightarrow A_\infty) \neq 0$ by Proposition. \square

Proof of Theorem. (\Rightarrow) part is easy (cf. [1, Proposition 2]). We assume that $\text{Ker}(N_{1,0}: A_1 \rightarrow A_0) \subseteq \text{Ker}(j_{1,\infty}: A_1 \rightarrow A_\infty)$. From the commutative diagram (cf. [2, Theorem 7])

$$(3) \quad \begin{array}{ccc} A_1 & \xrightarrow{\sim} & X/\nu_{0,1}Y \\ N_{1,0} \downarrow & & \downarrow \text{projection} \\ A_0 & \xrightarrow{\sim} & X/Y, \end{array}$$

we see that $\text{Ker}(N_{1,0}: A_1 \rightarrow A_0) = Y/\nu_{0,1}Y$. It follows from the assumption and Proposition that