

The Automorphism Group of the Klein Curve in the Mapping Class Group of Genus 3

By Atsushi MATSUURA

Department of Mathematical Sciences, University of Tokyo
(Communicated by Heisuke HIRONAKA, M. J. A., Sept. 12, 1996)

Let R be a compact Riemann surface of genus $g \geq 2$. Then $\text{Aut}(R)$, the automorphism group of R , can be embedded into the mapping class group (for its definition, see [1, Ch. 4]) or the Teichmüller group Γ_g of genus g ;

$$(1) \quad \iota : \text{Aut}(R) \hookrightarrow \Gamma_g \simeq \text{Out}^+(\pi_1(R)) = \text{Aut}^+(\pi_1(R))/\text{Int}(\pi_1(R)).$$

Here, $\text{Aut}^+(\pi_1(R))$ consists of the automorphisms of $\pi_1(R)$ inducing the trivial action on $H_2(\pi_1(R), \mathbf{Z}) \simeq \mathbf{Z}$.

Recall the Hurwitz theorem, which states that

$$(2) \quad \# \text{Aut}(R) \leq 84(g - 1).$$

If the equality holds in (2), then R is called a Hurwitz Riemann surface and $\text{Aut}(R)$ is called a Hurwitz group.

Let X be the Klein curve of genus 3 defined by the equation

$$x^3y + y^3z + z^3x = 0.$$

It is well known that X is a Hurwitz Riemann surface; $G := \text{Aut}(X)$ is isomorphic to $PSL_2(\mathbf{F}_7)$ and has order 168.

Now let us forget about the Klein curve, and consider an orientable compact C^∞ surface X of genus 3. We define the canonical generators of $\pi_1(X, b)$ with base point b as in the figure 1. They satisfy the fundamental relation

$$(3) \quad (\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1})(\alpha_2\beta_2\alpha_2^{-1}\beta_2^{-1})(\beta_3\alpha_3\beta_3^{-1}\alpha_3^{-1}) = 1.$$

Let $\tilde{\varphi}_2, \tilde{\varphi}_3, \tilde{\varphi}_7$ be the elements of $\text{Aut}^+(\pi_1(X))$ defined by

$$\begin{aligned} \tilde{\varphi}_2(\alpha_1) &= \alpha_2\beta_2^{-1}\alpha_2^{-1}\alpha_1^{-1}\beta_3^{-1}\beta_2 \\ \tilde{\varphi}_2(\beta_1) &= \beta_2^{-1}\beta_3\beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1} \\ \tilde{\varphi}_2(\alpha_2) &= \beta_3^{-1}\alpha_2^{-1} \\ \tilde{\varphi}_2(\beta_2) &= \alpha_2\beta_3\beta_2^{-1}\alpha_2^{-1} \\ \tilde{\varphi}_2(\alpha_3) &= \alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1^{-1}\alpha_1^{-1}\alpha_3\alpha_2^{-1} \\ \tilde{\varphi}_2(\beta_3) &= \alpha_2\beta_3\alpha_2^{-1}, \\ \tilde{\varphi}_3(\alpha_1) &= \alpha_2\beta_3\alpha_3^{-1}\alpha_1\alpha_2\beta_2\alpha_2^{-1} \\ \tilde{\varphi}_3(\beta_1) &= \alpha_2\beta_2^{-1}\alpha_2^{-1}\alpha_1^{-1}\alpha_3\alpha_1\alpha_2\beta_2\alpha_2^{-1} \\ \tilde{\varphi}_3(\alpha_2) &= \alpha_3^{-1}\alpha_1\beta_1\alpha_1^{-1} \\ \tilde{\varphi}_3(\beta_2) &= \alpha_1\beta_1^{-1}\alpha_1^{-1}\alpha_3\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1^{-1} \end{aligned}$$

$$\begin{aligned} \tilde{\varphi}_3(\alpha_3) &= \alpha_2\beta_2\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1 \\ \tilde{\varphi}_3(\beta_3) &= \alpha_1\beta_1^{-1}\alpha_1^{-1}\alpha_3\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1, \end{aligned}$$

$$\begin{aligned} \tilde{\varphi}_7(\alpha_1) &= \beta_1^{-1}\alpha_1^{-1}\alpha_3\beta_3^{-1}\alpha_2^{-1} \\ \tilde{\varphi}_7(\beta_1) &= \alpha_2\beta_3\alpha_3^{-1}\alpha_1\alpha_2\beta_2\alpha_2^{-1}\beta_3^{-1}\alpha_2^{-1} \\ \tilde{\varphi}_7(\alpha_2) &= \alpha_2\beta_2^{-1}\alpha_2^{-1}\alpha_1^{-1} \\ \tilde{\varphi}_7(\beta_2) &= \alpha_1\alpha_2\beta_2\beta_3\alpha_3^{-1} \\ \tilde{\varphi}_7(\alpha_3) &= \beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1}\alpha_3^{-1}\alpha_1\beta_1\alpha_1^{-1} \\ \tilde{\varphi}_7(\beta_3) &= \alpha_1\alpha_2\beta_2\alpha_3^{-1}\alpha_1\beta_1\alpha_1^{-1}. \end{aligned}$$

Then, we have the following:

Theorem 1. (1) *The classes φ_i of $\tilde{\varphi}_i$ in $\text{Out}^+(\pi_1(X))$ generate a subgroup H of Γ_3 , which is isomorphic to $PSL_2(\mathbf{F}_7)$.*

(2) *Moreover, if X is the Klein curve, then H is conjugate to the image of ι .*

Outline of the proof. (1) First note that $H \neq \{1\}$, because the action of H on the homology group $H_1(X, \mathbf{Z})$ is not trivial. By direct computation using (3), we have

$$(4) \quad \begin{aligned} \tilde{\varphi}_2^2 = \tilde{\varphi}_3^3 = \tilde{\varphi}_7^7 = 1, \quad \tilde{\varphi}_2\tilde{\varphi}_3\tilde{\varphi}_7 = 1, \\ (\tilde{\varphi}_7\tilde{\varphi}_3\tilde{\varphi}_2)^4 = [\text{conjugation by } \alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1]. \end{aligned}$$

For example,

$$\begin{aligned} \tilde{\varphi}_3^2 \cdot \beta_3 &= (\alpha_2^{-1}\beta_2\alpha_2\alpha_1\alpha_3\alpha_1^{-1}\alpha_2^{-1}\beta_2^{-1}\alpha_2)(\alpha_3\alpha_1^{-1}\beta_1^{-1}\alpha_1) \\ &\quad \times (\alpha_1^{-1}\beta_1\alpha_1\alpha_3^{-1}\alpha_2^{-1}\beta_2\alpha_2\beta_1^{-1}\alpha_1) \\ &\quad \quad \quad (\alpha_1^{-1}\beta_1\alpha_1\alpha_3^{-1}) \\ &\quad \times (\beta_1\alpha_2^{-1}\beta_2^{-1}\alpha_2\beta_2\alpha_2) \\ &\quad \quad \quad (\alpha_2^{-1}\beta_3^{-1}\alpha_3\alpha_1^{-1}\alpha_2^{-1}\beta_2^{-1}\alpha_2) \\ &\quad \times (\alpha_2^{-1}\beta_2\alpha_2\alpha_1\alpha_3^{-1}\alpha_1^{-1}\alpha_2^{-1}\beta_2^{-1}\alpha_2) \\ &\quad \quad \quad (\alpha_2^{-1}\beta_2\alpha_2\alpha_1\alpha_3^{-1}\beta_3\alpha_2) \\ &= \alpha_2^{-1}\beta_2\alpha_2(\alpha_1\beta_1\alpha_2^{-1}\beta_2^{-1}\alpha_2\beta_2\beta_3^{-1}\alpha_3^{-1}\beta_3)\alpha_2 \\ &= \alpha_2^{-1}\beta_2\alpha_2\beta_1\alpha_1\alpha_3^{-1}\alpha_2, \end{aligned}$$

hence

$$\begin{aligned} \tilde{\varphi}_3^3 \cdot \beta_3 &= (\alpha_3\alpha_1^{-1}\beta_1^{-1}\alpha_1)(\alpha_1^{-1}\beta_1\alpha_2^{-1}\beta_2^{-1}\alpha_2\alpha_3\alpha_1^{-1}\beta_1^{-1}\alpha_1) \\ &\quad \times (\alpha_1^{-1}\beta_1\alpha_1\alpha_3^{-1}) \\ &\quad \quad \quad (\alpha_2^{-1}\beta_2\alpha_2\alpha_1\alpha_3\alpha_1^{-1}\alpha_2^{-1}\beta_2^{-1}\alpha_2) \\ &\quad \times (\alpha_2^{-1}\beta_2\alpha_2\alpha_1\alpha_3^{-1}\beta_3\alpha_2) \\ &\quad \quad \quad (\alpha_2^{-1}\beta_2^{-1}\alpha_2^{-1}\beta_2\alpha_2\beta_1^{-1})(\alpha_1^{-1}\beta_1\alpha_1\alpha_3^{-1}) \\ &= \beta_3. \end{aligned}$$

From (4) we obtain

$$(5) \quad \varphi_2^2 = \varphi_3^3 = \varphi_7^7 = \varphi_2\varphi_3\varphi_7 = (\varphi_7\varphi_3\varphi_2)^4 = 1$$

in $\text{Out}^+(\pi_1(X))$. Since (5) is the presentation of