

## On Hasse Zeta Functions of Enveloping Algebras of Solvable Lie Algebras

By Takako FUKAYA

Department of Mathematics, Tokyo Institute of Technology  
(Communicated by Shokichi IYANAGA, M. J. A., Oct. 14, 1996)

**1. Introduction 1.1.** In [3], we generalized the definitions of Hasse zeta functions of commutative finitely generated rings over the ring  $\mathbf{Z}$  of integers, to non-commutative rings. In this paper we compute the Hasse zeta functions of the enveloping algebras of completely solvable Lie algebras having  $\mathfrak{p}$ -mappings.

For a (not necessarily commutative) finitely generated ring  $A$  over  $\mathbf{Z}$ , in [3] we defined the Hasse zeta function  $\zeta_A(s)$  of  $A$  by

$$\zeta_A(s) = \prod_{r \geq 1} \zeta_{A,r}(s)$$

where  $r$  runs over integers  $\geq 1$  and,

$$\zeta_{A,r}(s) = \prod_{\mathfrak{p}} \exp \sum_{n=1}^{\infty} \frac{\#\mathfrak{S}_{A,r}(\mathbf{F}_{\mathfrak{p}^n})}{n} (\mathfrak{p}^{-s})^n$$

where  $\mathfrak{S}_{A,r}$  is a certain scheme of finite type over  $\mathbf{Z}$ ,  $\mathfrak{p}$  runs over prime numbers, and  $\mathbf{F}_{\mathfrak{p}^n}$  is a finite field with  $\mathfrak{p}^n$  elements, so the function  $\zeta_{A,r}(s)$  coincides with the product of Weil's zeta functions of  $\mathfrak{S}_{A,r} \otimes_{\mathbf{Z}} \mathbf{F}_{\mathfrak{p}}$  [2] for all prime numbers  $\mathfrak{p}$ . We do not review the definition of  $\mathfrak{S}_{A,r}$ , but what we need in this paper is that for the algebraic closure  $K$  of  $\mathbf{F}_{\mathfrak{p}}$ ,  $\mathfrak{S}_{A,r}(K)$  is identified with the set of all isomorphism classes of  $r$ -dimensional irreducible representations of  $A$  over  $K$ , and  $\mathfrak{S}_{A,r}(\mathbf{F}_{\mathfrak{p}^n})$  is identified with the  $\text{Gal}(K/\mathbf{F}_{\mathfrak{p}^n})$ -fixed part of  $\mathfrak{S}_{A,r}(K)$ .

It has the expression

$$\zeta_A(s) = \prod_M (1 - N(M)^{-s})^{-1}$$

where  $M$  runs over the isomorphism classes of finite simple  $A$ -modules and  $N(M) = \#\text{End}_A(M)$ .

**1.2.** Recall that a solvable Lie algebra  $\mathfrak{g}$  over a field is said to be completely solvable if  $[\mathfrak{g}, \mathfrak{g}]$  is nilpotent. (See [1].)

We obtain the following result.

**Theorem 1.3.** *Let  $R$  be a commutative finitely generated ring over  $\mathbf{Z}$ . Let  $\mathfrak{g}$  be a Lie algebra over  $R$  which is free of finite rank  $n$  as an  $R$ -module, and let  $A$  be the universal enveloping algebra of  $\mathfrak{g}$ . Assume that for each maximal ideal  $\mathfrak{m}$  of  $R$ ,  $\mathfrak{g}/\mathfrak{m}\mathfrak{g}$  is a completely solvable Lie algebra over  $R/\mathfrak{m}$  and*

*has a  $\mathfrak{p}$ -mapping (see [1]). Then we have that the function  $\zeta_A(s)$  converges, and*

$$\zeta_A(s) = \zeta_R(s - n).$$

**Remark 1.3.1.** *For  $x \in \mathfrak{g}$ , let  $\text{ad}(x)$  be the inner derivation of  $\mathfrak{g}$  defined by  $x$ , that is,  $\text{ad}(x)(y) = [x, y]$  for  $y \in \mathfrak{g}$ . For a Lie algebra  $\mathfrak{g}$  over a field of characteristic  $\mathfrak{p}$ ,  $\mathfrak{g}$  has a  $\mathfrak{p}$ -mapping  $[\mathfrak{p}]$  if and only if the following condition holds: For any  $x \in \mathfrak{g}$ , there exists  $y \in \mathfrak{g}$  such that  $(\text{ad}(x))^{\mathfrak{p}} = \text{ad}(y)$ .*

**1.4. Example.** Every nilpotent Lie algebra  $\mathfrak{g}$  such that  $\mathfrak{g}^{\mathfrak{p}} = 0$  ( $\mathfrak{g}^i$  is defined by  $\mathfrak{g}^0 = \mathfrak{g}$  and  $[\mathfrak{g}^i, \mathfrak{g}] = \mathfrak{g}^{i+1}$  for  $i \geq 0$ ) satisfies the condition of Theorem 1.3. This is because  $(\text{ad}(x))^{\mathfrak{p}} = 0$  for any  $x \in \mathfrak{g}$ .

In section 2, we prove Theorem 1.3.

I would like to express my hearty gratitude to Professor Kazuya Kato for advice and encouragement.

**2. Proof of Theorem 1.3.** In this section we prove Theorem 1.3.

**2.1.** The zeta functions  $\zeta_R(s)$ ,  $\zeta_A(s)$  are products of  $\zeta_{R/\mathfrak{m}}(s)$ ,  $\zeta_{A/\mathfrak{m}A}(s)$  over all maximal ideals  $\mathfrak{m}$  of  $R$ , respectively, and  $A/\mathfrak{m}A$  are the universal enveloping algebras of  $\mathfrak{g}/\mathfrak{m}\mathfrak{g}$  over the finite fields  $R/\mathfrak{m}$ . So we may assume that  $R$  is a finite field  $k$  of characteristic  $\mathfrak{p}$ . Let  $K$  be the algebraic closure of  $k$ .

Theorem 1.3 follows from

**Proposition 2.2.** *Let  $\mathfrak{g}$  be a completely solvable Lie algebra over a finite field  $k$  of characteristic  $\mathfrak{p} > 0$  of finite dimension  $n$  which has a  $\mathfrak{p}$ -mapping  $[\mathfrak{p}]$ . Let  $A$  be the universal enveloping algebra of  $\mathfrak{g}$ , and let  $\mathbf{F}_q$  be a finite extension of  $k$ . Then we have*

$$\#\mathfrak{S}_A^k(\mathbf{F}_q) = q^n$$

*where  $\mathfrak{S}_A = \prod_{r \geq 1} \mathfrak{S}_{A,r}$  and  $\mathfrak{S}_A^k(\mathbf{F}_q)$  denotes the set of  $\mathbf{F}_q$ -rational points of  $\mathfrak{S}_A$  as a  $k$ -scheme.*

We prove Proposition 2.2 in 2.3 and 2.4.

**2.3.** There is a surjective map  $\varphi$  from  $\mathfrak{S}_A^k(K)$  onto  $K^{\oplus n}$ , the direct sum of  $n$  copies of  $K$ . Fix a basis  $(e_i)_{1 \leq i \leq n}$  of  $\mathfrak{g}$ . For an element  $x$  of