

## A Form of Classical Liouville Theorem

By Mitsuru NAKAI<sup>\*)†)</sup> and Toshimasa TADA<sup>\*\*)</sup>

(Communicated by Kiyosi ITÔ, M. J. A., Nov. 12, 1997)

The Liouville theorem in the theory of harmonic functions (cf. e.g. Axler *et al.* [1]) states that any nonnegative harmonic function  $u$  on the  $d$ -dimensional Euclidean space  $\mathbf{R}^d$  ( $d \geq 2$ ) reduces to a constant. It naturally occurs the question how much the condition for  $u$  to be nonnegative can be relaxed (see, e.g. Doob [3]). Recently Bourdon [2] proposed, among other related things, the following interesting generalization of the Liouville theorem :

**Theorem A** (Liouville Theorem). *If  $u$  is a harmonic function on  $\mathbf{R}^d$  and satisfies*

$$(1) \quad \liminf_{|x| \rightarrow \infty} \frac{u(x)}{|x|} \geq 0,$$

*then  $u$  is a constant function on  $\mathbf{R}^d$ .*

Bourdon gave an elementary and simple proof to the above result by using only the mean value property of harmonic functions originally due to an ingenious idea of Nelson [6](cf. also [1]). In contrast with the Liouville theorem in the theory of complex functions it is natural to consider Theorem A as a special case of the following result :

**Theorem B** (Liouville Theorem). *If  $u$  is a harmonic function on  $\mathbf{R}^d$  and satisfies*

$$(2) \quad \liminf_{|x| \rightarrow \infty} \frac{u(x)}{|x|^{n+1}} \geq 0$$

*for some nonnegative integer  $n$ , then  $u$  is a harmonic polynomial on  $\mathbf{R}^d$  of degree at most  $n$ .*

Clearly the  $n = 0$  case of Theorem B is nothing but Theorem A. The Bourdon proof of Theorem A seems not to be straightforwardly ap-

plied to that for Theorem B. It has been constantly our claim (cf. e.g. [4]) that the Fourier expansion method is one of the best tools to handle harmonic functions as far as their domains of definition are rotationally invariant such as  $\mathbf{R}^d$ . The purpose of this note is to give a proof to Theorem B by using the Fourier expansion, and actually, we prove Theorem B in the following superficially more general form :

**3. Theorem** (Liouville Theorem). *Suppose that  $u$  is a harmonic function on  $\mathbf{R}^d$  and that there exists an increasing divergent sequence  $(r_m)_{m \geq 1}$  of positive numbers  $r_m$  such that*

$$(4) \quad \liminf_{m \rightarrow \infty} \left( \min_{|x|=r_m} \frac{u(x)}{|x|^{n+1}} \right) \geq 0$$

*for some nonnegative integer  $n$ , then  $u$  is a harmonic polynomial on  $\mathbf{R}^d$  of degree at most  $n$ .*

*Proof.* We use the polar coordinate  $x = r\xi$  for points  $x \in \mathbf{R}^d$ , where  $r = |x| \geq 0$  and  $\xi = x/|x| \in S^{d-1}$  for  $x \neq 0$  and  $\xi = (1, 0, \dots, 0) \in S^{d-1}$  for  $x = 0$  for definitness. Here  $S^{d-1}$  is the unit sphere  $\{x \in \mathbf{R}^d : |x| = 1\}$ . We choose and then fix an orthonormal basis  $\{S_{kj} : j = 1, \dots, N(k)\}$  of the subspace of all spherical harmonics of degree  $k$  of  $L^2(S^{d-1}, d\sigma)$ , where  $d\sigma$  is the area element on  $S^{d-1}$ . Then  $\{S_{kj} : j = 1, \dots, N(k) ; k = 0, 1, \dots\}$  is a complete orthonormal system in  $L^2(S^{d-1}, d\sigma)$ . We have, as the special case of the addition theorem,

$$\sum_{j=1}^{N(k)} S_{kj}(\xi)^2 = \frac{N(k)}{\sigma_d},$$

where  $\sigma_d$  is the surface area  $\sigma(S^{d-1})$  of  $S^{d-1}$ . Here  $N(0) = 1$  and

$N(k) = (2k + d - 2)\Gamma(k + d - 2)/\Gamma(k + 1)\Gamma(d - 1)$  for  $k = 1, 2, \dots$ . For simplicity we set  $A_k := \sqrt{N(k)/\sigma_d}$  so that

$|S_{kj}(\xi)| \leq A_k$  ( $j = 1, \dots, N(k) ; k = 0, 1, \dots$ ). Then we have the following expansion of  $u(r\xi)$  in terms of spherical harmonics  $\{S_{kj}\}$  :

$$(5) \quad u(r\xi) = \sum_{k=0}^{\infty} \left( \sum_{j=1}^{N(k)} a_{kj} S_{kj}(\xi) \right) r^k,$$

where  $a_{kj}$  ( $j = 1, \dots, N(k) ; k = 0, 1, \dots$ ) are

<sup>\*)</sup> Department of Mathematics, Nagoya Institute of Technology.

<sup>\*\*)</sup> Department of Mathematics, Daido Institute of Technology.

<sup>†)</sup> Present address: Department of Mathematics, Daido Institute of Technology.

1991 Mathematics Subject Classification. Primary 31B05.

This work was partly supported by Grant-in-Aid for Scientific Research, Nos. 08640194, 08640243, Ministry of Education, Science, Sports and Culture, Japan.