

On an Infinite Family of Elliptic Curves with Rank ≥ 14 over \mathbf{Q}

By Shoichi KIHARA

Department of Neuropsychiatry School of Medicine Tokushima University

(Communicated by Shokichi IYANAGA, M. J. A., Feb. 12, 1997)

In [2], Nagao constructed an example \mathcal{E} of elliptic curve over $\mathbf{Q}(t)$ with rank ≥ 13 . In this paper, we show in utilizing \mathcal{E} and a method introduced in our previous paper [3] that there are infinitely many elliptic curves with rank ≥ 14 over \mathbf{Q} .

The curve given in [2] was

$$\mathcal{E}: y^2 = (9s^2 + 211950)x^4 + (-2700s^2 - 63901710)x^3 + (-18s^4 + 396150s^2 + 6706476489)x^2 + (2700s^4 - 29575350s^2 - 284435346600)x + 9s^6 - 159200s^4 + 891699592s^2 + 4156297690000$$

where $s = (-t^2 + 23550)/(2t)$. As was shown in [2], there are following 13 points on \mathcal{E} .

$$\begin{aligned} P_1 &= (s + 148, 662s^2 + 66873s + 1868944), \\ P_2 &= (s + 116, -554s^2 - 39687s - 191632), \\ P_3 &= (s + 104, -526s^2 - 28497s + 163372), \\ P_4 &= (s + 57, 508s^2 - 19332s - 368809), \\ P_5 &= (s + 25, 580s^2 - 49116s + 566825), \\ P_6 &= (s, -670s^2 + 69759s - 2038700), \\ P_7 &= (-s + 148, -662s^2 + 66873s - 1868944), \\ P_8 &= (-s + 116, 554s^2 - 39687s + 191632), \\ P_9 &= (-s + 104, 526s^2 - 28497s - 163372), \\ P_{10} &= (-s + 57, -508s^2 - 19332s + 368809), \\ P_{11} &= (-s + 25, -580s^2 - 49116s - 566825), \\ P_{12} &= (-s, 670s^2 + 69759s + 2038700), \\ P_{13} &= ((s + 703)/15, (-224s^3 - 844s^2 + 900484s + 2161725)/75). \end{aligned}$$

Next, let us consider the following elliptic curve:

$$C: q^2 = p(p - 13728)(p + 80472).$$

$(-27456, -7742592)$ is on C , and it is easy to see that this point is of infinite order in the Mordell-Weil group of C , so that C has positive rank.

Let $\mathbf{Q}(C)$ be the function field of C . Now we consider \mathcal{E} over $\mathbf{Q}(C)$, like in [3], by specializing

$$t = q/(2p).$$

Then we have the point $P_{14} = (x_{14}, y_{14})$ on \mathcal{E} , where

$$\begin{aligned} x_{14} &= (-1104719616 - p^2 + 708q)/(4q) \\ y_{14} &= (240419869705111928832 - 12282065003400192p + 1177306772832p^2 + 11117812p^3 + 197p^4 - 108850233203712q - 98532p^2q)/(4q^2). \end{aligned}$$

Theorem 1. P_1, \dots, P_{14} are independent points.

Proof. This is shown by specializing $(p, q) = (-27456, -7742592)$. Let R_1, \dots, R_{14} be the rational points obtained from P_1, \dots, P_{14} by the above specialization. By using calculation system PARI, we see that the determinant of the matrix $(\langle R_i, R_j \rangle)$ ($1 \leq i, j \leq 14$) associated to the canonical height is 2344685535688581.87. Since this determinant is non-zero, we see that R_1, \dots, R_{14} are independent points.

So we see that P_1, \dots, P_{14} are independent.

Q.E.D.

Now by the theorem 20.3 in [1], specializing (p, q) to elements of the Mordell-Weil group of C , we have

Theorem 2. There are infinitely many elliptic curves over \mathbf{Q} with rank ≥ 14 .

References

- [1] J. H. Silverman: The arithmetic of elliptic curves. Graduate Texts in Math., vol. 106, Springer-Verlag, New York (1986).
- [2] K. Nagao: An example of elliptic curve over $\mathbf{Q}(T)$ with rank ≥ 13 . Proc. Japan Acad., **70A**, 152–153 (1994).
- [3] S. Kihara: On the rank of the elliptic curve $y^2 = x^3 + k$. II. Proc. Japan Acad., **72A**, 228–229 (1996).