

## A Theta Product Formula for Jackson Integrals Associated with Root Systems

By Masahiko ITO

Graduate School of Polymathematics, Nagoya University

(Communicated by Heisuke HIRONAKA, M. J. A., April 14, 1997)

### Jackson integrals associated with root systems.

Let  $\mathfrak{a}$  be an  $n$ -dimensional vector space over  $\mathbf{R}$  with an inner product  $\langle \cdot, \cdot \rangle$ . Let  $R \subset \mathfrak{a}^*$  be an irreducible reduced root system and  $W_R$  be the group generated by orthogonal reflections with respect to the hyperplane perpendicular to  $\alpha \in R$ , the so-called *Weyl group* associated with  $R$ . Let  $P$  be the *weight lattice* of  $R$  defined by  $\{\mu \in \mathfrak{a}^*; \langle \mu, \alpha^\vee \rangle \in \mathbf{Z} \text{ for any } \alpha \in R\}$ , where  $\alpha^\vee = 2\alpha / \langle \alpha, \alpha \rangle$ . We fix a base  $\{\alpha_1, \dots, \alpha_n\} \subset R$  and its fundamental weights  $\{\chi_1, \dots, \chi_n\} \subset P; \langle \chi_i, \alpha_j^\vee \rangle = \delta_{ij}$ . The inner product and the reflections are uniquely extended linearly to  $\mathfrak{h} = \mathbf{C} \otimes_{\mathbf{R}} \mathfrak{a}$ . We sometimes identify the vector space  $\mathfrak{h}$  with its dual  $\mathfrak{h}^*$  via the inner product  $\langle \cdot, \cdot \rangle: \mu(\alpha) = \langle \mu, \alpha \rangle$ .

Let  $\bar{X}$  be an algebraic torus of dimension  $n$ , isomorphic to  $(\mathbf{C}^\times)^n$ . We can embed  $P$  in  $\bar{X}$  by the mapping

$$\begin{aligned} \mathfrak{h}^* &\rightarrow \bar{X}; \chi = \nu_1 \chi_1 + \dots + \nu_n \chi_n \\ &\mapsto q^\chi := (q^{\nu_1}, \dots, q^{\nu_n}) \end{aligned}$$

where  $q = e^{2\pi\sqrt{-1}\tau}$ ,  $\text{Im}\tau > 0$ . We denote by  $X$  the lattice subgroup  $\{(q^{\nu_1}, \dots, q^{\nu_n}); \nu_i \in \mathbf{Z} (i = 1, \dots, n)\} \subset \bar{X}$ . We identify  $P$  with  $X$ . Each  $\alpha \in \mathfrak{h}^*$  defines a monomial  $t^\alpha := t_1^{\langle \chi_1, \alpha^\vee \rangle} \dots t_n^{\langle \chi_n, \alpha^\vee \rangle}$  for  $t = (t_1, \dots, t_n) \in \bar{X}$ . To each  $\alpha \in R$ , let  $k_\alpha$  be a complex number such that  $k_\alpha = k_\beta$  if  $|\alpha| = |\beta|$ .

We introduce the following function of  $t = (t_1, \dots, t_n)$  on  $\bar{X}$  (see [3]):

$$\Phi_R(k; t) = t^\ell \prod_{\alpha > 0} \frac{(q^{1-k_\alpha} t^\alpha)_\infty}{(q^{k_\alpha} t^\alpha)_\infty}$$

where  $(x)_\infty = \prod_{\nu=1}^\infty (1 - xq^\nu)$ ,  $\ell = \frac{1}{2} \sum_{\alpha > 0} (1 - 2k_\alpha)$  and “ $\alpha > 0$ ” means  $\alpha$  is a positive root of  $R$ . For simplicity we sometimes abbreviate  $\Phi_R(k; t)$  by  $\Phi_R(t)$ . The function  $\Phi_R(k; t)$  is quasi-symmetric with respect to  $W_R$ :

$$\sigma \Phi_R(k; t) = \Phi_R(k; \sigma^{-1}(t)) = U_\sigma(t) \cdot \Phi_R(k; t), \quad \sigma \in W_R$$

where  $U_\sigma(t)$  is a *pseudo-constant*, i.e. a  $q$ -periodic function with respect to  $t \in \bar{X}$  such that

$$U_\sigma(t) = \prod_{\substack{\alpha > 0 \\ \sigma\alpha < 0}} t^{(2k_\alpha - 1)\alpha} \frac{\vartheta(q^{k_\alpha} t^\alpha)}{\vartheta(q^{1-k_\alpha} t^\alpha)}$$

for the Jacobi elliptic theta function  $\vartheta(x) = (x)_\infty (q/x)_\infty (q)_\infty$ .  $\{U_\sigma(t)\}_{\sigma \in W_R}$  satisfies the *one cocycle* condition such that  $U_{\sigma\sigma'}(t) = U_\sigma(t) \cdot \sigma U_{\sigma'}(t)$ .

We let  $\Delta_R$  denote the *Weyl denominator* defined by  $\Delta_R(t) := \prod_{\alpha > 0} (t^{\frac{\alpha}{2}} - t^{-\frac{\alpha}{2}})$ . Let us define  $\Phi'_R(k; t) := \Phi_R(k; t) \cdot (-1)^{\frac{|R|}{2}} \Delta_R(t)$ . Then, the function  $\Phi'_R(k; t)$  also has the *quasi-symmetry*  $\sigma \Phi'_R(k; t) = \text{sgn}\sigma \cdot U_\sigma(t) \cdot \Phi'_R(k; t)$ ,  $\sigma \in W_R$ .

**Definition.** We now consider the *Jackson integral associated with  $R$*  defined by

$$\begin{aligned} J_R(k; \xi) &:= \int_{\{0, \xi_\infty\}_q} \Phi'_R(k; t) \frac{d_q t_1}{t_1} \wedge \dots \wedge \frac{d_q t_n}{t_n} \\ &= (1 - q)^n \sum_{x \in X} \Phi'_R(k; q^x \xi) \end{aligned}$$

where  $\xi = (\xi_1, \dots, \xi_n)$  is an arbitrary point of  $\bar{X}$  and  $q^x \xi$  means  $(q^{\nu_1} \xi_1, \dots, q^{\nu_n} \xi_n)$ .

It is obvious that the Jackson integral  $J_R(k; \xi)$  is a  $q$ -periodic function of  $\xi \in \bar{X}$  if it is convergent:

$$J_R(k; q^x \xi) = J_R(k; \xi).$$

Let  $\Gamma_q(x)$  denote the  $q$ -gamma function  $(1 - q)^{1-x} (q)_\infty / (q^x)_\infty$ .

**Conjecture** (product formula). *The Jackson integral  $J_R(k; \xi)$  can be expressed as follows:*

$$(1) \quad J_R(k; \xi) = \prod_{\alpha > 0} \frac{\Gamma_q(1 - \langle \rho_k, \alpha^\vee \rangle) \Gamma_q(-\langle \rho_k, \alpha^\vee \rangle)}{\Gamma_q(1 - k_\alpha - \langle \rho_k, \alpha^\vee \rangle) \Gamma_q(k_\alpha - \langle \rho_k, \alpha^\vee \rangle + \delta_\alpha) \frac{\xi^{-k_\alpha \alpha} (\xi^\alpha)}{\vartheta(q^{k_\alpha} \xi^\alpha)}}$$

up to a positive integer, where  $\delta_\alpha = 1$  if  $\alpha$  is a simple root,  $\delta_\alpha = 0$  otherwise, and  $\rho_k = \frac{1}{2} \sum_{\alpha > 0} k_\alpha \alpha$ .

**Proposition.** *The Jackson integrals of  $A_n$ -type,  $B_2$ -type and  $G_2$ -type have the following formulae:*

$$J_{A_n}(\beta; \xi) = (n + 1) \prod_{j=1}^n$$