

Simplified proof of an order preserving operator inequality

By Takayuki FURUTA

Department of Applied Mathematics, Science University of Tokyo,
1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601
(Communicated by Kiyosi ITÔ, M. J. A., Sept. 14, 1998)

A capital letter means a bounded linear operator on a Hilbert space.

Theorem 1. *If $A \geq B \geq 0$ with $A > 0$, then for $1 \geq q \geq t \geq 0$ and $p \geq q$,*
(1) $A^{q-t+r} \geq \{A^{r/2}(A^{-t/2}B^pA^{-t/2})^s A^{t/2}\}^{q/[(p-t)s+t]}$
for $s \geq 1$ and $r \geq t$.

We cite the following results to prove a simplified proof of Theorem 1 in [3].

Theorem A [1]. *If $A \geq B \geq 0$, then for each $r \geq 0$*
(A-1) $(A^{r/2}A^pA^{r/2})^{1/q} \geq (A^{r/2}B^pA^{r/2})^{1/q}$
holds for $p \geq 0$ and $q \geq 1$
with $(1+r)q \geq p+r$.

The domain drawn for p, q and r in Figure is the best possible one [4] for (A-1). Theorem A implies Löwner-Heinz inequality :

(#) $A \geq B \geq 0$ ensures $A^\alpha \geq B^\alpha$ for $\alpha \in [0, 1]$.

Lemma [2]. *Let X be a positive invertible and Y be an invertible. For any real number λ ,*
 $(YXY^*)^\lambda = YX^{1/2}(X^{1/2}Y^*YX^{1/2})^{\lambda-1}X^{1/2}Y^*$.

Proof of Theorem 1. First of all, we prove that if $A \geq B \geq 0$ with $A > 0$, then
(*) $A^q \geq \{A^{t/2}(A^{-t/2}B^pA^{-t/2})^s A^{t/2}\}^{q/[(p-t)s+t]}$ for $1 \geq q \geq t \geq 0, p \geq q$ and $s \geq 1$.

In case $2 \geq s \geq 1$, as $s-1, q/[(p-t)s+t] \in [0, 1]$ and $A^t \geq B^t$ by (#), we have (2) by Lemma and (#)

$$\begin{aligned} (2) \quad B_1 &= \{A^{t/2}(A^{-t/2}B^pA^{-t/2})^s A^{t/2}\}^{q/[(p-t)s+t]} \\ &= \{B^{p/2}(B^{p/2}A^{-t}B^{p/2})^{s-1}B^{p/2}\}^{q/[(p-t)s+t]} \\ &\leq \{B^{p/2}(B^{p/2}B^{-t}B^{p/2})^{s-1}B^{p/2}\}^{q/[(p-t)s+t]} = \\ &B^q \leq A^q = A_1 \text{ by (#)} \end{aligned}$$

for $1 \geq q \geq t \geq 0, p \geq q$ and $2 \geq s \geq 1$. Repeating (2) for $A_1 \geq B_1 \geq 0$, then we have


$$(3) \quad A_1^{q_1} \geq \{A_1^{t_1/2}(A_1^{-t_1/2}B_1^{p_1}A_1^{-t_1/2})^{s_1}A_1^{t_1/2}\}^{q_1/[(p_1-t_1)s_1+t_1]}$$

for $1 \geq q_1 \geq t_1 \geq 0, p_1 \geq q_1$ and $2 \geq s_1 \geq 1$. Put $1 = q_1 \geq t_1 = t/q \geq 0$ and

$$\begin{aligned} p_1 &= [(p-t)s+t]/q \geq q_1 = 1 \text{ in (3). Then} \\ (4) \quad A^q &\geq \{A^{t/2}[A^{-t/2}A^{t/2}(A^{-t/2}B^pA^{-t/2})^s A^{t/2}]^{s_1}A^{t/2}\}^{q/[(p-t)s_1+t]} \\ &= \{A^{t/2}(A^{-t/2}B^pA^{-t/2})^{ss_1}A^{t/2}\}^{q/[(p-t)s_1+t]} \text{ for } 1 \geq q \geq t \\ &\geq 0, p \geq q \text{ and } 4 \geq ss_1 \geq 1. \end{aligned}$$

Repeating this process from (2) to (4), we obtain

(*) for $1 \geq q \geq t \geq 0, p \geq q$ and any $s \geq 1$. Put $A_2 = A^q$ and $B_2 = \{A^{t/2}(A^{-t/2}B^pA^{-t/2})^s A^{t/2}\}^{q/[(p-t)s+t]}$ in (*). Then $A_2 \geq B_2 \geq 0$ by (*), so by Theorem A we have
(5) $A_2^{1+r_2} \geq (A_2^{r_2/2}B_2^{p_2}A_2^{r_2/2})^{(1+r_2)/(p_2+r_2)}$ for $p_2 \geq 1$ and $r_2 \geq 0$.
We have only to put $r_2 = (r-t)/q \geq 0$ and $p_2 = [(p-t)s+t]/q \geq 1$ in (5) to obtain (1).



References

- [1] T. Furuta: $A \geq B \geq 0$ assures $(B^rA^pB^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0, p \geq 0, q \geq 1$ with $(1+2r)q \geq p+2r$. Proc. Amer. Math. Soc., **101**, 85-88 (1987).
- [2] T. Furuta: Extension of the Furuta inequality and Ando-Hiai log majorization. Linear Alg and Its Appl., **219**, 139-155 (1995).
- [3] T. Furuta, T. Yamazaki, and M. Yanagida: Order preserving operator inequalities via Furuta inequality (preprint).
- [4] K. Tanahashi: Best possibility of the Furuta inequality. Proc. Amer. Math. Soc., **124**, 141-146 (1996).