194. On Axiom Systems of Propositional Calculi. XXIV

By Kiyoshi Iséki
Kobe University

(Comm. by Kinjirô Kunugi, M.J.A., Oct. 12, 1966)

In his well known book [2], A. Church defines a propositional calculus P_1 which is equivalent to the classical propositional calculus. The calculus with a propositional constant 0 (which corresponds to the false proposition) is given by the following axioms:

1 $CpCqp$,
2 $CCpCqrCCpqCpr$,
3 $CCCp00p$.

In [1], by two inference rules of substitution and detachment we proved that the first two axioms imply the following theses:

4 Cpp,
5 $CCpqCCqrCpr$,
6 $CCqrCCpqCpr$,
7 $CCpCqrCqCpr$,
8 $CCpCpqCpq$.

In this note, using substitution and detachment rules, we shall show that the calculus is the classical one. To do so, put $Np=Cp0$, then the axiom 3 means $CNNpp$. We deduce an axiom system by J. Lukasiewicz (see [3]):

a $CCpqCCqrCpr$,
b $CCNppp$,
c $CpCNpq$.

The thesis a follows from the axioms 1 and 2, as already well-known so we must prove that the theses b and c hold in the P_1-calculus. For the proofs, we use the proofline method.

1 $p/CCCp00p$, $q/0$ $*C3$—9,
9 $C0CCCp00p$.
2 $p/0$, $q/CCp00$, r/p $*C9$—$C1$ $p/0$, $q/Cp0$—10,
10 $C0p$.
1 $p/C0q$, q/p $*C10$ p/q—11,
11 $CpC0q$.
2 $q/0$, r/q $*C10$—12,
12 $CCp0Cpq$.
6 $p/Cp0$, q/p, r/q $*C12$—13,
13 $CpCCp0q$.

This means $CpCNpq$, which is the thesis c.

7 $p/Cp0$, q/p, $r/0$ $*C4$ $p/Cp0$—14,