

74. On the Dimension of an Orbit-space

By Takashi KARUBE

Faculty of Engineering, Gifu University

(Comm. by K. KUNUGI, M.J.A., June 12, 1961)

Let G be a locally compact transformation group satisfying the second axiom of countability and acting on a locally compact Hausdorff space M , and H be a compact invariant subgroup of G . Then in a natural way the set of all orbits under H becomes a locally compact Hausdorff space, which is called "the orbit-space of M under H " and denoted by $D(M; H)$, and the factor group $G^* = G/H$ acts on $D(M; H)$ as a transformation group (cf. [4], p. 61). In this note we prove that

$$\dim G(x) = \dim H(x) + \dim D(G(x); H) \quad \text{for } x \in M. \quad (\text{A})$$

This is a generalization of a result obtained by Montgomery and Zippin ([5], p. 783, cf. Corollary of the present note). If $G(x)$ is finite dimensional, then $D(G(x); H)$ is locally the topological product of a Euclidean cube by a zero dimensional set closed in $D(G(x); H)$ (cf. Karube [3]); so that the equation (A) gives us the almost complete knowledge about the local topology of such an orbit-space as the above.

We now prove the equation (A).

1) *Let G be finite dimensional.* Let p be the natural projection of M onto $D(M; H)$, and \tilde{x} the image of the point x under p . Let π be the natural mapping of G onto G^* , F^* the group of all elements of G^* leaving the point \tilde{x} fixed, and F the complete inverse image of F^* under π . It is easy to see that $F(x) = H(x)$ and $G_x = F_x$ where G_x and F_x are stability subgroups of the point x . By the theorems of Yamanoshita [6] we have

$$\begin{aligned} \dim G &= \dim F + \dim G/F, \\ \dim G &= \dim G(x) + \dim G_x, \\ \dim F &= \dim F(x) + \dim F_x = \dim H(x) + \dim G_x, \\ \dim G/F &= \dim G^*/F^* = \dim G^*(\tilde{x}) = \dim D(G(x); H). \end{aligned}$$

Since G_x is finite dimensional, we have (A).

2) *Let $G(x)$ be finite dimensional.* There exists an open subgroup G' of G such that G'/G_0 is compact where G_0 is the identity component of G . Since $G'(x)$ is finite dimensional, G' is effectively finite dimensional on $G'(x)$. In fact, there must be a connected compact invariant subgroup K' of G' which is idle on $G'(x)$ and such that G'/K' is finite dimensional (cf. [3]). Let G'_1 be the factor group G'/K' , ρ the natural mapping of G' onto G'_1 , H' the intersection of H and G' , and H'_1 the image of H' under ρ . Since G'_1 is finite dimensional we have