

On the Representation Spaces of the Jordanian Plane

Natalia K Iyudu*

Department of Pure Mathematics, Queen's University Belfast, Belfast BT7 1NN, UK

We announce here a number of results concerning representation theory of the algebra $R = k \langle x, y \rangle / (xy - yx - y^2)$, known as Jordanian plane (or Jordan algebra). We consider the question on 'classification' of finite-dimensional modules over the Jordan algebra. Complete description of irreducible components of the representation variety $\text{mod}(R, n)$ is given for any dimension n . It is obtained on the basis of the stratification of this variety related to the Jordan normal form of Y . Any irreducible component of the representation variety contains only one stratum related to a certain partition of n and is the closure of this stratum. The number of irreducible components therefore is equal to the number of partitions of n .

As a preparation for the above result we describe the complete set of pairwise non-isomorphic irreducible modules S_a over the Jordan algebra, and the rule how they could be glued to indecomposables. Namely, we show that $\text{Ext}^1(S_a, S_b) = 0$, if $a \neq b$.

We study then properties of the image algebras in the endomorphism ring. Particularly, images of representations from the most important stratum, corresponding to the full Jordan block Y . This stratum turns out to be the only building block for the analogue of the Krull-Remark-Schmidt decomposition theorem on the level of irreducible components. Along this line we establish an analogue of the Gerstenhaber-Taussky-Motzkin theorem on the dimension of algebras generated by two commuting matrices. Another fact concerns with the tame-wild question for those image algebras. We show that all image algebras of n -dimensional representations are tame for $n \leq 4$ and wild for $n \geq 5$.

We consider here the quadratic algebra given by the presentation $R = k \langle x, y \rangle / (xy - yx - y^2)$. This algebra appears in various different contexts in mathematics and physics. First of all, it is a kind of quantum plane: one of the two Auslander regular algebras of global dimension two, as it is mentioned in the Artin and Shelter paper [1]. The other one is the usual quantum plane $k \langle x, y \rangle / (xy - qyx)$. It served as one of basic examples for the foundation of noncommutative geometry [2]. There were also studies of deformations of $GL(2)$ analogous to $GL_q(2)$ with respect to R in 80-90th in Manin's 'Quantum group' [3,4], where this algebra appeared under the name Jordan algebra.

This algebra is also a simplest element in the class of RIT (relativistic internal time) algebras. The latter was defined and investigated [5-9]. The class of RIT algebras arises from a modification of the Poincare algebra of the Lorentz group $SO(3,1)$ by means of introducing the additional generator corresponding to the relativistic internal time. The algebra R above is a RIT algebra of type (1,1). Our studies of this algebra are partially reported "Classification of finite dimensional representations of one noncommutative quadratic algebra" [10].

Let us mention that R is a subalgebra of the first Weyl algebra A_1 . The latter has no finite dimensional representations, but R turns out to have quite a rich structure of them. Category of finite dimensional modules over R contains, for example, as a full subcategory $\text{mod } GP(n, 2)$, where $GP(n, 2)$ is the Gelfand-Ponomarev algebra [11] with the nilpotency degrees of variables x and y , n and 2 respectively. On the other hand we show that R is residually finite dimensional.

We are interested here in representations over an algebraically closed field k of characteristic 0. Sometimes we just suppose $k = \mathbb{C}$, this will be pointed out separately. We denote throughout the category of all R -modules by $\text{Mod } R$, the category of finite dimensional R -modules by $\text{mod } R$ and ρ_n stands for an n -dimensional representation of R .

The most important question on finite dimensional representations one can ask, is to classify them. This question may be solved directly by parametrization of isoclasses of indecomposable modules, for example, for tame algebras. But when the algebra is wild the problem turns to a description of orbits under GL_n action by simultaneous conjugation in the space $\text{mod}(R, n)$ of n -dimensional representations. One can think of the latter space also as of a variety of tuples of $n \times n$ matrices (corresponding to generators) satisfying the defining relations of the algebra R .

It is commonly understood that the first step in the study of this variety should be the description of its irreducible components. This approach leads to such famous results of this kind as Kashiwara-Saito [12] description of the irreducible components of Lusztig's nilpotent variety via the crystal basis [13].

We show that irreducible components of the representation space $\text{mod}(R, n)$ of the Jordan algebra could be completely described for any dimension n .

The first key point for this description is the choice of a stratification of the Jordan variety. We choose a stratum \mathcal{U}_p corresponding to the partition $\mathcal{P} = (n_1, \dots, n_k)$, $n = n_1 + \dots + n_k$, $n_1 \geq \dots \geq n_k$, consisting of the pairs of matrices (X, Y) , where Y has the Jordan normal form defined by \mathcal{P} and X, Y satisfy the defining relation.

Theorem 12.1 Any irreducible component K_i of the representation variety $\text{mod}(R, n)$ of the Jordan algebra contains only one stratum \mathcal{U}_p from the stratification related to the Jordan normal form of Y , and is the closure of this stratum.

The number of irreducible components of the variety $\text{mod}(R, n)$ is equal to the number of the partitions of n .

The variety $\text{mod}(R, n)$ is equidimensional, the dimension of components is n^2 .

The importance of examples of algebras for which the irreducible components of $\text{mod}(R, n)$ could be described for each n was emphasized [14] and it is mentioned there that known cases are restricted to algebras of finite representation type (i.e., there are only finitely many

*Corresponding author: Natalia K Iyudu, Department of Pure Mathematics, Queen's University Belfast, Belfast BT7 1NN, UK, E-mail: n.iyudu@qub.ac.uk

Received July 21, 2015; Accepted August 03, 2015; Published August 31, 2015

Citation: Iyudu NK (2015) On the Representation Spaces of the Jordanian Plane. J Generalized Lie Theory Appl S1: e001. doi:10.4172/1736-4337.S1-e001

Copyright: © 2015 Iyudu NK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.