



ANNALES DE L'INSTITUT HENRI POINCARÉ

PROBABILITÉS ET STATISTIQUES

Limit law of a second class particle in TASEP with non-random initial condition	<i>P. L. Ferrari, P. Ghosal and P. Nejjar</i>	1203–1225
Homogenization theory for the random conductance model with degenerate ergodic weights and unbounded-range jumps	<i>F. Flegel, M. Heida and M. Slowik</i>	1226–1257
Activated random walk on a cycle	<i>R. Basu, S. Ganguly, C. Hoffman and J. Ricbey</i>	1258–1277
Non-Gaussian limit theorem for non-linear Langevin equations driven by Lévy noise	<i>A. Kulik and I. Pavlyukevich</i>	1278–1315
Estimating functions for SDE driven by stable Lévy processes	<i>E. Clément and A. Gloter</i>	1316–1348
Inverting the cut-tree transform	<i>L. Addario-Berry, D. Dieuleveut and C. Goldschmidt</i>	1349–1376
A large deviation principle for empirical measures on Polish spaces: Application to singular Gibbs measures on manifolds	<i>D. García-Zelada</i>	1377–1401
Edge of spiked beta ensembles, stochastic Airy semigroups and reflected Brownian motions	<i>P. Y. G. Lamarre and M. Sbkolnikov</i>	1402–1438
Pathwise differentiability of reflected diffusions in convex polyhedral domains	<i>D. Lipsbutz and K. Ramanan</i>	1439–1476
Einstein relation and linear response in one-dimensional Mott variable-range hopping	<i>A. Faggionato, N. Gantert and M. Salvi</i>	1477–1508
Limit laws for self-loops and multiple edges in the configuration model	<i>O. Angel, R. van der Hofstad and C. Holmgren</i>	1509–1530
Gravitation versus Brownian motion	<i>S. Banerjee, K. Burdzy and M. Duarte</i>	1531–1565
On the roughness of the paths of RBM in a wedge	<i>P. Lakner, J. Reed and B. Zwart</i>	1566–1598
Diffusion limit for the random walk Metropolis algorithm out of stationarity	<i>J. Kuntz, M. Ottobre and A. M. Stuart</i>	1599–1648
Mixing times for a constrained Ising process on the two-dimensional torus at low density	<i>N. S. Pillai and A. Smith</i>	1649–1678
Quasi-independence for nodal lines	<i>A. Rivera and H. Vanneuville</i>	1679–1711
Liouville quantum gravity spheres as matings of finite-diameter trees	<i>J. Miller and S. Sheffield</i>	1712–1750
Active phase for activated random walks on \mathbb{Z}^d, $d \geq 3$, with density less than one and arbitrary sleeping rate	<i>L. Taggi</i>	1751–1764
Supremum estimates for degenerate, quasilinear stochastic partial differential equations	<i>K. Dareiotis and B. Gess</i>	1765–1796
On thin local sets of the Gaussian free field	<i>A. Sepúlveda</i>	1797–1813