

on general mechanics has already reached its fourth edition. The second and third editions were issued under the superintendence of the author. Dr. Wien has had the advantage of the author's manuscript notes, found after his death, in preparing the volume for the fourth edition. No changes have, however, been made beyond the correction of printer's errors and the removal of slight obscurities. One of the best testimonies that we can give to the care and ability which Kirchhoff devoted to the original volume is to say that those who possess any one of the first three issues will not find it necessary to buy this last edition.

ERNEST W. BROWN.

*Leçons élémentaires sur la théorie des formes et ses applications géométriques*, a l'usage des candidats a l'agregation des sciences mathématiques. Par H. ANDOYER. Paris, Gauthier-Villars et Fils, 1898. 4to, 184 pp. Lithographed.

THIS title represents accurately the contents of the book. A restricted list of topics is adequately treated. Geometrical applications are plentiful but do not encroach upon nor obscure the purely algebraic theory. The treatment is perhaps elementary, but in style simplicity is less noticeable than brevity. The work as a whole is more nearly a syllabus than a textbook for the beginner, and as a syllabus it cannot fail to become widely known and valued.

Binary and ternary forms are introduced, but of binary forms only linear, bilinear, quadric, cubic, and quartic forms are discussed; and of ternary, only linear, bilinear, and quadric. The bilinear ternary form in cogredient variables I do not remember having seen in any earlier text, although Salmon's *Higher plane curves* gives a brief geometrical discussion of skew reciprocity; its inclusion here together with the form bilinear in contragredient variables must be commended by geometers. Duality as a method stands certainly on a par with projectivity.

Two points of excellence are worthy of special mention. The first is that from the outset stem forms are assumed to contain several sets of variables; and this convention is observed in the case of each particular form, an unlimited number of sets of cogredient variables being adjoined. Of course this amplifies the complete system of each stem form, but the extension is easy, since no new types occur. The second novel merit is that the term *polar* is so defined as to include such operations as Aronhold's. If so-called variables are replaced by cogredient variables, the process is ordinarily called a polar operation, and M. Andoyer sees no