

not seem to apply when $n > 2$. This suggests the important question: Can any fundamental circle group (discontinuous ζ -group that is representable with real coefficients) be defined arithmetically as a Γ_n for a sufficiently high value of n ?

UNIVERSITY OF ILLINOIS,
December, 1909.

A NEW ANALYTICAL EXPRESSION FOR THE NUMBER π , AND SOME HISTORICAL CONSIDERATIONS.

BY DR. G. VACCA.

IN the common exposition of the history of mathematics, more attention is given to results than to methods, and it is only rarely that old theorems and demonstrations are translated into the modern and living mathematical language. The great mathematicians of the eighteenth century (Euler, Lagrange, and others) have made important contributions to this difficult work. It seems to me that even to-day something can be done in this direction, and many new results can be obtained by a careful reading of the ancient classics of mathematics. The following is an example:

Let $f(x)$ be a function defined by the relation

$$f(x) = \frac{1}{2}(x + |x|)$$

and suppose that $f^n(x)$ means the result of the operation f applied n times to the number x . Then we have, if $i = \sqrt{-1}$,

$$(1) \quad \frac{2}{\pi} = \lim_{n \rightarrow \infty} f^n(i).$$

This elegant theorem can be easily proved. It gives a possible analytical definition of the number π .

It is now to be observed that this *new* formula is only the analytical expression (using the geometrical representation of complex quantities of Gauss) of a series of points (approaching the vertex) of the quadratrix (*τετραγωνίζουσα*) of Deinostratos.*

We may try to transform the second member of (1) into a real expression. This can be done by elementary methods. But the result is nothing else than the well-known infinite product of complicated quadratic radicals first given by Vieta.†

* Pappi Coll. Math., lib. IV, prop. 25.

† Vieta, Opera, ed. Schooten, Lugd. Batav., 1646, p. 400.