

## LINEAR OPERATIONS ON FUNCTIONS OF BOUNDED VARIATION

T. H. HILDEBRANDT

The object of this note is to give a form for the most general linear continuous operation on the space of functions of bounded variation on a finite interval, say  $0 \leq x \leq 1$ , the norm of the space being the total variation.

This form is obtained by setting up an equivalent space. For this purpose let  $\mathfrak{I}$  be the class of elements  $I$  consisting of any finite number of non-overlapping intervals  $i_1, \dots, i_n$  of the interval  $(0, 1)$ . If  $(x_p, y_p)$  are endpoints of  $i_p$ , define the function of interval sets  $\beta(I) = \sum_{p=1}^n [\alpha(y_p) - \alpha(x_p)]$  corresponding to the function  $\alpha(x)$  of bounded variation. Then  $\beta(I)$  is a bounded function on  $\mathfrak{I}$ . Define  $\|\beta\|$  in the usual way as the least upper bound of  $|\beta(I)|$  for  $I$  on  $\mathfrak{I}$ . Then the space  $\mathfrak{B}$  of additive set functions  $\beta$  thus normed is equivalent to the space  $\mathfrak{A}$  of functions  $\alpha(x)$  of bounded variation with  $\|\alpha\| = V\alpha = \int_0^1 |d\alpha|$ ,\* for obviously  $\|\beta\| \leq \|\alpha\| \leq 2\|\beta\|$ . Further, if  $\alpha_1$  corresponds to  $\beta_1$  and  $\alpha_2$  to  $\beta_2$ , then  $\beta_1 + \beta_2$  corresponds to  $\alpha_1 + \alpha_2$  and  $c\beta$  to  $c\alpha$ , and conversely.

It is now an easy matter to determine the most general linear functional operation on the space  $\mathfrak{B}$ . Following the lines of reasoning of my paper *On bounded linear functional operations*,† one finds that for any linear continuous operation  $L$  on the space  $\mathfrak{B}$  there exists an additive function  $\gamma$  of sets  $E$  of elements  $I$ , such that  $L(\beta) = \int \beta d\gamma$ , the integral being of the  $L$  or  $S$  type as defined in the paper quoted, and extended over the class of all subsets of elements of  $\mathfrak{I}$ . Because of the relationship between the functions  $\beta$  and  $\alpha$  this gives the most general linear operation in the space  $\mathfrak{A}$ .

It might be noted that a similar reasoning applies to the set of interval functions  $\alpha(i)$  where  $\sum_{p=1}^n \alpha(i_p) = \beta(I)$  is a bounded function on  $\mathfrak{I}$ ; or, more generally, that a similar result holds in the space of bounded functions on a general range, with norm the least upper bound of the absolute value of the function on the range.

UNIVERSITY OF MICHIGAN

\* Note that in the space  $\mathfrak{A}$  two functions for which  $V(\alpha_1 - \alpha_2) = \int |d(\alpha_1 - \alpha_2)| = 0$  are regarded as equivalent. To obtain uniqueness, the condition  $\alpha(0) = 0$  can be added. If we wish that  $\|\alpha\| = 0$  imply  $\alpha = 0$  for all  $x$ , we may choose  $\|\alpha\| = |\alpha(0)| + V\alpha$ . The space  $\mathfrak{B}_1$  corresponding is defined by  $\beta_1(I) = \alpha(0) + \sum_{p=1}^n [\alpha(y_p) - \alpha(x_p)] = \alpha(0) + \beta(I)$  and  $\|\beta_1(I)\| = |\alpha(0)| + \|\beta(I)\|$ . Reasoning similar to the above can be carried through in this case also.

† Transactions of this Society, vol. 36 (1934), pp. 868-875.