

REPRESENTATIONS OF BOOLEAN ALGEBRAS

ORRIN FRINK, JR.

There are several proofs in the literature¹ of M. H. Stone's theorem on the representation of Boolean algebras by sets [2, 4, 5, 7, 8, 9]. This note contains a simplified version of Stone's original proof, adapted to the following set, I-IV, of postulates for a Boolean algebra B in terms of the special element 0 and the undefined operations *product* ab and *negation* b' . It is assumed that 0 is in B , and that if a , b , and c are in B , then ab and b' are in B , and

I. $ab = ba$.

II. $a(bc) = (ab)c$.

III. $aa = a$.

IV. $ab = a$ if and only if $ab' = 0$.

Replacing b by a in IV gives V: $aa' = 0$. Since $a0 = a(aa') = (aa)a' = aa' = 0$, we have VI: $a0 = 0$.

DEFINITIONS. A *point* is a set P of elements of B such that

α . The element 0 is not in P .

β . If a is in P and b is in P , then ab is in P .

γ . P is maximal with respect to properties α and β .

The set R_a of all points P which contain a is defined to be the *representative set* corresponding to the element a .

LEMMA 1. If ab is in P , then a is in P .

PROOF. If a were not in P , then P would not be maximal, since a and all products pa , where p is in P , could then be added to P without disturbing α , since if $pa = 0$, then $pab = 0$.

LEMMA 2. If a is not equal to 0, then a is in some point P .

PROOF. All sets of elements of B which contain a and satisfy α and β form a system S partially ordered by set inclusion. Any linearly ordered subsystem L of S has an upper bound in S , namely the union of all members of L . Hence by Zorn's lemma [10, 11], there exists in S at least one maximal element P .

THEOREM. The correspondence between elements a of B and their representative sets R_a is an isomorphism; that is, 1. $R_{ab} = R_a \cap R_b$; 2. $R_{a'} = C(R_a)$; 3. if $R_a = R_b$, then $a = b$.

¹ See also N. H. McCoy and D. Montgomery, Duke Mathematical Journal, vol. 3 (1937), pp. 455-459.