

## CIRCUMSCRIBED CUBES IN EUCLIDEAN $n$ -SPACE

BY S. S. CAIRNS

Communicated by R. H. Bing, May 20, 1959

Let  $E^n$  be a euclidean  $n$ -space with a rectangular cartesian coordinate system  $(x) = (x_1, \dots, x_n)$ , and let  $(y)$  be any system which is a rotation of  $(x)$ . Let  $A \subset E^n$  be a closed bounded set containing  $n+1$  linearly independent points. Its circumscribed  $(y)$ -box is the set  $a_i \leq y_i \leq b_i$  ( $i=1, \dots, n$ ) where  $a_i$  and  $b_i$  are the respective minimum and maximum values of  $y_i$  on  $A$ . Let  $c_i = b_i - a_i$  be interpreted as a function on the space  $R_{n-1}$  of rotations of coordinate systems, which is also the rotation space of the unit  $(n-1)$ -sphere  $S^{n-1} \subset E^n$ .

Let  $f: R_{n-1} \rightarrow E^n$  be the function which maps  $r \in R_{n-1}$  onto the point  $(c_1(r), \dots, c_n(r))$ , relative to the fixed initial coordinate system  $(x)$ . Let  $D$  be the diagonal  $x_1 = \dots = x_n$  in  $E^n$ . The circumscribed  $(y)$ -box corresponding to a point  $r \in R_{n-1}$  is an  $n$ -cube if and only if  $f(r) \in D$ . Accordingly,  $K = f^{-1}(D)$ , a subspace of  $R_{n-1}$ , will be called the *space of circumscribed  $n$ -cubes* of  $A$ . Its structure can be studied by means of the mapping  $f$ . For the purpose of this study the significant properties are as follows: (1)  $f$  is a continuous mapping of  $R_{n-1}$  into the region  $x_i > 0$  ( $i=1, \dots, n$ ) of  $E^n$  (2)  $f(R_{n-1})$  is symmetric with respect to  $D$ . This second property follows from the fact that all possible permutations of axial directions can be achieved in a symmetric way through rotations. There is no need to distinguish between the two possible senses on a given  $y_i$ -direction, since the value of  $c_i$  is the same for both. Hence, one gets odd as well as even permutations of the  $c$ 's.

Let  $T^{n-1}$  be the simplex in  $E^n$  with vertices at the unit points on the  $(x)$ -axes. A central projection from the origin carries the mapping  $f$  into a continuous mapping  $g: R_{n-1} \rightarrow T^{n-1}$  where  $g(R_{n-1})$  is symmetric in the barycentric coordinates on  $T^{n-1}$ . The inverse image  $g^{-1}(q)$ , where  $q$  is the barycenter of  $T^{n-1}$ , is identical with  $f^{-1}(D) = K$ . This leads to the following result.

**THEOREM.** *The space of circumscribed cubes of a closed subset of euclidean  $n$ -space containing  $n+1$  independent points is the inverse image  $K = g^{-1}(q)$  of the center of an  $(n-1)$ -simplex  $T^{n-1}$  under a continuous mapping  $g: R_{n-1} \rightarrow T^{n-1}$ , where  $R_{n-1}$  is the rotation space of an  $(n-1)$ -sphere and where  $g(R_{n-1})$  is symmetric in the barycentric coordinates on  $T^{n-1}$ .*

Any particular circumscribed  $n$ -cube is the  $(y)$ -cube for a system  $(y)$  obtainable from  $(x)$  without rotating any axis by more than