

REPRESENTING MEASURES FOR POINTS IN A UNIFORM ALGEBRA

BY ERRETT BISHOP

Communicated by Walter Rudin, September 9, 1963

Recent years have seen much effort put into attempts to develop an "abstract" function theory. In the complex domain, this has led to the study of the structure of uniform algebras. A uniform algebra \mathfrak{A} is a family of continuous complex-valued functions on a compact Hausdorff space X , which contains the function 1, which is closed with respect to the algebraic operations of addition and multiplication by complex scalars, which is topologically closed in the uniform norm, and which distinguishes points of X . Succinctly, \mathfrak{A} is a closed separating unitary subalgebra of the Banach algebra $C(X)$ of all continuous complex-valued functions on X . Standard examples are obtained by taking X to be a compact subset of complex Euclidean space C^n and \mathfrak{A} to be the closed subalgebra of $C(X)$ generated by the constants and the coordinate functions on C^n .

Some of the most suggestive ideas in the theory of uniform algebras come from a paper of Gleason [1]. He calls elements x and y in X *equivalent* if

$$d(x, y) = \sup\{ |f(x) - f(y)| : f \in \mathfrak{A}, \|f\| \leq 1 \} < 2.$$

This is an equivalence relation on X . The equivalence classes are called *parts* or *Gleason parts*. The parts have various other characterizations, obtainable from the elementary conformal geometry of the disc $D = \{z : |z| \leq 1\}$. We shall need to know, in particular, that if there exists a sequence $\{h_n\}$ of elements of \mathfrak{A} , $\|h_n\| \leq 1$, $|h_n(x)| \rightarrow 1$ as $n \rightarrow \infty$, then $|h_n(y)| \rightarrow 1$ as $n \rightarrow \infty$ if x and y are in the same part.

Gleason showed, in a special case, that parts can be given a certain characterization in terms of representing measures. A representing measure μ_x for a point x in X is a non-negative Baire measure on X such that $\int f d\mu_x = f(x)$ for all f in \mathfrak{A} . The following theorem generalizes Gleason's result to the general situation.

THEOREM 1. *If x and y are in the same part, there exists $c > 0$ and representing measures μ_x for x and μ_y for y such that $c\mu_x \leq \mu_y$ and $c\mu_y \leq \mu_x$.*

PROOF. Let $C_r(X)$ be the Banach space of all continuous real-valued functions on X . Let R be the set of all f in $C_r(X)$ such that $f + ig \in \mathfrak{A}$ for some g in $C_r(X)$. Let c be a constant, $0 < c < 1$, such that