

A LOCALLY COMPACT SEPARABLE METRIC SPACE IS ALMOST INVARIANT UNDER A CLOSED MAPPING

BY EDWIN DUDA

Communicated by P. E. Conner, October 25, 1963

For a given mapping (continuous transformation) f of a topological space X onto a topological space Y it has always been of interest to determine what properties of X carry over to Y . Under the hypothesis that f is a closed mapping it is known that normality [1], and paracompactness [5] are invariants. If X is metric and f is closed then as a consequence of results of Vainštejn [2], Whyburn [1], and Stone [4], it is known that Y is weakly separable if and only if each point inverse has a compact frontier. From [1] we obtain the result that if X is perfectly separable, f is closed and Y is weakly separable then Y is a separable metric space. Let f be a closed mapping of a locally compact separable metric space X onto a topological space Y , or, equivalently, let G be an upper semi-continuous decomposition of X into closed sets. We say a set S is a *scattered* set if every subset of S is closed. Our results show that Y (or the decomposition space M) minus a scattered set S , which has at most a countable number of points, is also a locally compact separable metric space. The techniques of proof are standard and will not be included here.

THEOREM 1. *Let G be an upper semi-continuous decomposition of a locally compact separable metric space into closed sets. Let F be the union of the noncompact elements of G , M the decomposition space determined by G , and ϕ the natural mapping of X onto M . The following are valid.*

- (i) F is a closed set.
- (ii) For an arbitrary compact set K only a finite number of elements of G in F can intersect K .
- (iii) F contains at most countably many elements of G .
- (iv) The union of any subcollection of elements of G in F is a closed set.
- (v) M is weakly separable at y if and only if the frontier of $\phi^{-1}(y)$ is compact (Stone [4]).
- (vi) If $\{g_n\}$ is a convergent sequence of compact elements of G with a nonempty limiting set h , then the set $K = \bigcup g_n \cup h$ is compact.

Let F' be the subset of F composed of the union of the elements