

# ON THE THICKNESS OF THE COMPLETE GRAPH<sup>1</sup>

BY LOWELL W. BEINEKE AND FRANK HARARY

Communicated by J. W. T. Youngs, January 22, 1964

The *thickness*  $t(K_p)$  of the complete graph  $K_p$  with  $p$  points is the minimum number of planar subgraphs whose union is  $K_p$ . The purpose of this note is to outline a result which determines  $t(K_p)$  for four of every six consecutive integers  $p$ . A complete proof of this result will be published elsewhere.

THEOREM. *If  $p \equiv -1, 0, 1, 2 \pmod{6}$ , then*

$$(1) \quad t(K_p) = \left\lceil \frac{p+7}{6} \right\rceil.$$

In proving this theorem, we prescribe a labelling of  $n+1$  plane graphs, for any positive integer  $n$ . All the graphs contain the same  $6n+2$  points, but are constructed so that no two have a common line. Two of the points will be denoted by  $v$  and  $v'$ , and the others as  $u_k, v_k, w_k, u'_k, v'_k, w'_k$  for  $k=0, 1, \dots, n-1$ . All but one of the graphs are of the type indicated in Figure 1, where each of the six numbered triangles in  $G_k$  contains  $n-1$  other points and  $3(n-1)$  lines so that its interior is isomorphic with graph  $H$ .

The points of the  $n$  graphs  $G_k$  are labelled using an  $n \times n$  matrix  $A = (a_{ij})$ , whose entries are residue classes modulo  $n$ , where

$$(2) \quad a_{ij} = \left( (-1)^i \left\lfloor \frac{i}{2} \right\rfloor + (-1)^j \left\lfloor \frac{j}{2} \right\rfloor \right) \pmod{n}$$

with  $\lfloor x \rfloor$  indicating the greatest integer function as usual. We remark that one of the important properties of  $A$  is that each residue class appears exactly once in each row and each column.

The  $n-1$  points inside triangle  $u'_k v_k w'_k$  of graph  $G_k$  are labelled using the column, say the  $j$ th, whose first entry is  $a_{1j} = k$  as follows: if  $a_{ij} = h$ , the  $(i-1)$ st point down from  $v_k$  is labelled  $v_h$  or  $v'_h$  according as  $\min \{i, j\}$  is odd or even. The points inside triangle  $v_k u'_k w_k$  are similarly labelled, using  $u'_h$  and  $u_h$  instead of  $v_h$  and  $v'_h$  respectively. The points inside the other triangles are also labelled analogously.

Now, in the union of these  $n$  labelled graphs  $G_k$ , aside from  $v$  and  $v'$ , each point is adjacent with all but one of the other points. More-

---

<sup>1</sup> This research was supported by National Science Foundation grant GP-207.