

MIXED BOUNDARY VALUE PROBLEMS FOR NONSTRICT HYPERBOLIC EQUATIONS

BY RICHARD BEALS¹

Communicated by Felix Browder, December 29, 1971

The Cauchy problem for a class of nonstrict hyperbolic equations has been studied by Ohya and Leray [3], [5]. Recently Chazarain [2] has considered certain mixed problems for some fourth order nonstrict equations in the context of distribution semigroups. In this note we deal with mixed problems for a general class of (possibly) nonstrict equations of higher order, including those in [2]. The class of hyperbolic operators does not contain nor is it contained in those of [3], [5]. We do not require Gevrey class coefficients, nor that the principal part has real characteristics of constant multiplicity and therefore (see [4]) is a product of powers of strictly hyperbolic operators. We do, however, require the operators to be elliptic in the space variables alone and that the coefficients be independent of time. In this case recent results on the abstract Cauchy problem [1] can be applied. For simplicity we consider a single equation with smooth coefficients on a bounded region in euclidean space. Proofs will appear elsewhere.

Let $\Omega \subset \mathbf{R}^n$, $n \geq 2$, be a bounded open set with boundary $\partial\Omega$ a regularly imbedded C^∞ submanifold of \mathbf{R}^n . Let $a(y; \xi, \lambda)$ be the polynomial

$$\sum_{|\alpha|+j \leq p} a_{\alpha j}(y) \xi^\alpha \lambda^j, \quad y \in \bar{\Omega}, \xi \in \mathbf{C}^n, \lambda \in \mathbf{C}.$$

Let $a_0(y; \xi, \lambda)$ be the principal part, the terms of total degree p . We assume that, for $y \in \bar{\Omega}$, $\xi \in \mathbf{R}^n$, the roots of $a_0(y; \xi, \cdot)$ are all real and nonzero. This implies $p = 2r$ is even. Let $b_k(y; \xi, \lambda)$ be polynomials of order $p_k < p$, $k = 1, 2, \dots, r$, with principal part $b_{k0}(y; \xi, \lambda)$, $y \in \partial\Omega$, $\xi \in \mathbf{C}^n$, $\lambda \in \mathbf{C}$. We assume the coefficients of these polynomials are C^∞ . We also assume the appropriate version of the "root covering condition" familiar from elliptic boundary value problems. Specifically, for $y \in \partial\Omega$, let $\nu = \nu(y)$ be the unit vector in \mathbf{R}^n in the direction of the inner normal to $\partial\Omega$ at y , and let $W = W(y)$ be the subspace of \mathbf{R}^n orthogonal to ν . The assumptions imply that for $\xi \in W$ and $\text{Re } \lambda > 0$, or $\xi \in W \setminus \{0\}$ and $\lambda = 0$, $a_0(y; \mu\nu + i\xi, \lambda)$ has precisely r roots $\mu_1(\xi, \lambda), \dots, \mu_r(\xi, \lambda)$ with negative real part. Let

$$q(\mu; \xi, \lambda) = \Pi(\mu - \mu_j(\xi, \lambda)).$$

AMS 1970 subject classifications. Primary 35L35.

Key words and phrases. Mixed boundary value problems, hyperbolic equations, multiple characteristics.

¹ This research was supported in part by NSF Grant GP-28148.