

OPERATORS WITH DISCONNECTED SPECTRA ARE DENSE

BY DOMINGO A. HERRERO¹ AND NORBERTO SALINAS²

Communicated by Fred Gehring, December 13, 1971

ABSTRACT. It is proven that the set of all (bounded linear) operators on a complex infinite dimensional Banach space having disconnected spectra is an open uniformly dense subset of the algebra of all operators.

In [3, Problem 8], P. R. Halmos asked whether the set of all reducible operators in a complex infinite dimensional separable Hilbert space \mathcal{H} is uniformly dense in the algebra $\mathcal{L}(\mathcal{H})$ of all (bounded linear) operators on \mathcal{H} . In the present note we answer affirmatively a related question:

Is the set of all operators on a Banach space X having nontrivial complementary hyperinvariant subspaces dense in $\mathcal{L}(X)$? (Recall that a subspace \mathcal{M} of X is *hyperinvariant* for $T \in \mathcal{L}(X)$ if $A\mathcal{M} \subset \mathcal{M}$, for all $A \in \mathcal{L}(X)$ commuting with T [1]. Here and in what follows, *subspace* means *closed linear manifold*.)

Moreover, we proved the following stronger (see [4]) result:

THEOREM. *Let X be a complex infinite dimensional Banach space and let $T \in \mathcal{L}(X)$. Then, given any $\varepsilon > 0$, there exists an $A \in \mathcal{L}(X)$ such that (1) $\text{rank}(A) = 1$; (2) $\|A\| < \varepsilon$, and (3) the spectrum of $T + A$ is disconnected.*

PROOF. Let $\sigma(T)$ ($E(T)$, resp.) denote the spectrum (essential spectrum, resp.) of T .

Let λ_0 be any point of $E(T)$ such that $\text{Re } \lambda_0 = \max\{\text{Re } \lambda; \lambda \in E(T)\}$. Then, for every compact operator K , $\lambda_0 \in E(T + K) = E(T) \subset \sigma(T + K)$, and it follows from [4, Theorem 1] that, if there exists a $\lambda \in \sigma(T + K)$ such that $\text{Re } \lambda > \text{Re } \lambda_0$, then $\sigma(T + K)$ is disconnected, λ is an isolated point of $\sigma(T + K)$ such that $(T + K - \lambda)^n X$ is closed for every $n \geq 0$ and, if $\mathcal{M} = \bigcap_{n=1}^{\infty} (T + K - \lambda)^n X$ and $\mathcal{N} = \text{closure}\{\bigcup_{n=1}^{\infty} \ker(T + K - \lambda)^n\}$, then $\dim \mathcal{N} = \dim(X/\mathcal{M}) < \infty$.

Therefore, to complete the proof, it suffices to find an A satisfying (1), (2) and such that $\lambda_0 + \gamma \in \sigma(T + A)$ for some γ , $0 < \gamma < \varepsilon/2$.

Since $\lambda_0 \in \text{bdry } \sigma(T)$, there exists an $x \in X$ such that $\|x\| = 1$ and $\|(T - \lambda_0)x\| < \varepsilon/2$ (see [2, Chapter 7]). By Hahn-Banach theorem, there

AMS 1970 subject classifications. Primary 47A15; Secondary 47A10, 47A55.

Key words and phrases. Complementary subspaces, hyperinvariant subspaces, disconnected spectrum, essential spectrum.

¹ State University of New York at Albany, Albany, New York 12203. Research supported by NSF Grant GU3171.

² Institute for Advanced Study, Princeton, New Jersey 08540. Research supported in part by NSF Grant GP 7952X3.