

NONSEPARATING FUNCTION ALGEBRAS

BY L. Q. EIFLER

Communicated by Felix Browder, December 29, 1971

Let A be a function algebra on X (compact). We say A is a separating algebra on X if for each closed subset S of X and for each $x \in X \setminus S$ there exists f in A such that $f(x) = 0$ and f does not vanish on S . We say that A is essential on X if for each open subset U of X there is a continuous function $f \notin A$ such that f vanishes on X/U . Csordas and Reiter asked [2] if there exists a nonseparating, essential algebra A on a (connected) space X for which X is the maximal ideal space of A and also the Šilov boundary of A . We give an example of such an algebra and simple examples of nonseparating algebras.

Given a compact subset K of C^n , let $P(K)$ denote the uniform closure in $C(K)$ of the polynomials in z_1, \dots, z_n . An easy application of Hurwitz' theorem [1, p. 176] shows that the first three of the following algebras are nonseparating.

EXAMPLE 1. Let $\Delta = \{z: |z| \leq 1\}$. Then $P(\Delta \times \Delta)$ is nonseparating since $f(\Delta \times \Delta) = f(\{(z, w): |z| = 1 \text{ or } |w| = 1\})$ for each f in $P(\Delta \times \Delta)$. Also, $\Delta \times \Delta$ is the maximal ideal space of $P(\Delta \times \Delta)$.

EXAMPLE 2. The algebra $P([0, 1] \times \Delta)$ is nonseparating since $f([0, 1] \times \Delta) = f(\{(t, z): t = 0 \text{ or } |z| = 1\})$. Also, $[0, 1] \times \Delta$ is the maximal ideal space of $P([0, 1] \times \Delta)$.

EXAMPLE 3. If A is a separating algebra on $X = M(A)$ and if B is a function algebra on X containing A , then B is separating on X but not necessarily separating on $M(B)$. Let B denote the uniform closure in $C(\Delta)$ of polynomials in z and $|z|$. Then $P(\Delta) \subseteq B \subseteq C(\Delta)$. One can embed B into $P([0, 1] \times \Delta)$ by setting $F(r, z) = f(rz)$ for $0 \leq r \leq 1$ and $|z| \leq 1$. Now one can see that the maximal ideal space of B is $[0, 1] \times \Delta / \{0\} \times \Delta$ and so B is nonseparating on its maximal ideal space.

EXAMPLE 4. Let $S^2 = C \cup \{\infty\}$ and let $D = \{z: |z| < 1\}$. Let

$$B = \{f \in C(S^2): f \text{ is analytic in } D\}.$$

Then $M(B) = S^2$ and $f(S^2) = f(S^2 \setminus D)$ for each f in B . One easily checks that S^2 is the maximal ideal space of B . Fix f in B and assume f does not vanish on $S^2 \setminus D$. If f has a zero in D , then there exist $\alpha_1, \dots, \alpha_n \in D$ such that $g(z) = f(z) \prod_{k=1}^n z/(z - \alpha_k)$ belongs to B and vanishes only at 0. Given $\infty \geq r > 0$, define $\Gamma_r(\theta) = gr e^{i\theta}$. All of the curves Γ_r are homotopic

AMS 1970 subject classifications. Primary 46J10.
Key words and phrases. Function algebras.