

## ZEROS OF SUCCESSIVE DERIVATIVES OF ENTIRE FUNCTIONS

BY R. P. BOAS, JR. AND A. R. REDDY

Communicated by R. C. Buck, June 5, 1972

Let  $f(z)$  be a transcendental entire function. If  $r_k$  is the radius of the largest disk with center at 0 in which  $f^{(k)}(z)$  is zero-free, it is known that, when  $f(z)$  is of positive finite order  $\rho$  and  $\alpha > \rho$ , there is an infinite increasing sequence of values of  $k$  such that  $r_k \geq k^{(1/\alpha)-1}$  (Ålander [1] for  $\rho < 1$ ; stated by Pólya [4] for  $\rho > 1$  also; the first published proof for  $\rho > 1$  was given by Erdős and Rényi [3], where Ålander's result is misquoted as being for  $\rho > 1$ ). When  $\rho = 1$  and  $f(z)$  is of exponential type  $\tau$  it is known more precisely that  $r_k \geq c(\tau)$  (Takenaka [5]; for modern results see Buckholtz and Frank [2]).

We have established the existence of larger zero-free disks if they are no longer required to be centered at 0. Our principal results are as follows.

**THEOREM 1.** *If  $f(z)$  is an entire function at most of order 2, finite type, there is an arbitrarily large disk, somewhere in the plane, in which an infinity of  $f^{(k)}(z)$  are zero-free.*

This is a corollary of Ålander's theorem for  $\rho < 1$ , but not for  $1 \leq \rho \leq 2$ .

The conclusion of Theorem 1 fails for entire functions of order greater than 2.

**THEOREM 2.** *If  $\rho > 2$ , there is an entire function of order  $\rho$  such that, for some positive  $A$ , every disk, anywhere in the plane, of radius  $A$  contains a zero of every  $f^{(k)}(z)$ .*

**THEOREM 3.** *If  $f(z)$  is an entire function of finite order  $\rho \geq 2$ , and  $\alpha > \rho$ , there is a point  $z_0$  such that, for an infinity of  $k$ , we have  $f^{(k)}(z) \neq 0$  in  $|z - z_0| < k^{(1/\alpha)-1/2}$ .*

Theorem 3 shows that when we do not require the concentric zero-free disks to be centered at a prescribed point, they can be appreciably larger than in Pólya's theorem.

**THEOREM 4.** *If  $f(z)$  is an entire function, for every (arbitrarily large)  $c > 0$ , there is a  $z_0$  such that  $f^{(k)}(z) \neq 0$  in  $|z - z_0| < ck^{-1/2}$  for an infinity of  $k$ .*

**THEOREM 5.** *If  $f(z)$  is analytic in  $|z| < R$ , there are a (possibly small)  $c > 0$  and a point  $z_0$  in  $|z| < R$  such that  $f^{(k)}(z) \neq 0$  in  $|z - z_0| < ck^{-1/2}$  for an infinity of  $k$ .*

*AMS (MOS) subject classifications* (1970). Primary 30A64, 30A66; Secondary 30A08.

Copyright © American Mathematical Society 1973