

A CHARACTERIZATION OF INNER PRODUCT SPACES

BY R. A. TAPIA

Communicated by R. Creighton Buck, November 6, 1972

Let X be a real normed linear space and f a functional on X . Recall that by the first one-sided variation of f at x in the direction h we mean

$$f'_+(x)(h) = \lim_{t \rightarrow 0^+} \frac{[f(x + th) - f(x)]}{t}$$

and by the second one-sided variation of f at x in the directions h_1 and h_2 we mean

$$f''_+(x)(h_1, h_2) = \lim_{t \rightarrow 0^+} \frac{[f'_+(x + th_1)(h_2) - f'_+(x)(h_2)]}{t}.$$

Let

$$f(x) = \frac{1}{2}\|x\|^2 \quad \text{and} \quad (x, y) = f'_+(x)(y).$$

PROPOSITION 1. *Every normed linear space resembles an inner product space in the sense that*

- (i) (x, y) is well defined;
- (ii) $(x, x) \geq 0$ with equality if and only if $x = 0$;
- (iii) $\|x\| = (x, x)^{1/2}$.

Moreover, if X is an inner product space with inner product $[\cdot, \cdot]$, then $(\cdot, \cdot) = [\cdot, \cdot]$.

PROPOSITION 2. *The following are equivalent:*

- (i) X is an inner product space;
- (ii) (x, y) is symmetric;
- (iii) (x, y) is linear in the first variable;
- (iv) $f''_+(0)(x, y)$ is linear in the first variable.

The proofs of these and related results will appear elsewhere.

REFERENCES

1. J. P. Aubin, *Optimal approximation and characterization of the error and stability functions in Banach spaces*, J. Approximation Theory **3** (1970), 430–444. MR **43** #795.
2. A. Beurling and A. E. Livingston, *A theorem on duality mappings in Banach spaces*, Ark. Mat. **4** (1961), 405–411. MR **26** #2851.
3. R. Bonic and F. Reis, *A characterization of Hilbert space*, An. Acad. Brasil. Ci. **38** (1966), 239–241. MR **36** #1961.
4. F. E. Browder, *On a theorem of Beurling and Livingston*, Canad. J. Math. **17** (1965), 367–372. MR **31** #595.

AMS (MOS) subject classifications (1970). Primary 46B99, 46C10.

Key words and phrases. Inner product, differentiability of the norm.