

FIXED POINTS OF DISK ACTIONS

BY ROBERT OLIVER

Communicated October 30, 1975

As a sequel to a previous announcement [3], the author can now give a complete classification up to homotopy type of which spaces can occur as fixed point sets of smooth actions of a given compact Lie group on disks. The result is contained in Theorems 1 to 3 below. For a group G , G_0 denotes its identity component.

THEOREM 1. *Let G be a compact Lie group, and F a finite CW complex. Then there exists a smooth action of G on a disk with fixed point set having the homotopy type of F if and only if:*

1. $G \cong T^n$ ($n \geq 1$): F is \mathbf{Z} -acyclic;
 2. G_0 a torus and $|G/G_0| = p^a$ (p prime, $a \geq 1$): F is \mathbf{Z}_p -acyclic,
 3. G_0 not a torus or G/G_0 not of prime power order: $\chi(F) \equiv 1 \pmod{n_G}$
- for some fixed integer n_G .

In order to describe the calculations of n_G , some classes of finite groups are defined, as in [3] and [4]. G^1 denotes the class of all G with normal subgroup P of prime power order, such that G/P is cyclic. For q prime, G^q denotes the class of all G with normal subgroup $H \in G^1$ of q -power index. Then one gets

- THEOREM 2.**
1. *If G_0 is not a torus, then $n_G = 1$.*
 2. *If G_0 is a torus, then $n_G = n_{G/G_0}$.*
 3. *If G is finite, then $n_G = 0$ if and only if $G \in G^1$; if $G \notin G^1$ then for any prime q , $q \mid n_G$ if and only if $G \in G^q$.*

In Theorem 1, the necessity of the conditions in (1) and (2) follow from standard Smith theory. Sufficiency follows in (2) from Jones [2], and in (1) is trivial ($G * F$ is contractible and can be thickened up to a disk action by Theorem 6 of [4]).

For finite G , the existence of n_G and the calculations in Theorem 2, part 3, were proven in [4]. Furthermore, if G_0 is a torus and $G \supseteq G_0$, then F clearly has the homotopy type of the fixed point set of a disk action of G if and only if it does the same for G/G_0 , so $n_G = n_{G/G_0}$. The case where G_0 is nontoral will be dealt with below; the above theorems say that *any* finite homotopy type can occur as fixed point set for such G .

The following result, completing the calculation of n_G , was obtained in

AMS (MOS) subject classifications (1970). Primary 57E25; Secondary 55C35.

Copyright © 1976, American Mathematical Society